

# HEALTHY AGING

Presented by

CONTINUING PSYCHOLOGY EDUCATION INC.

7 CONTINUING EDUCATION CONTACT HOURS

“From a medical and public perspective, extending the period of successful aging, where people have sufficient cognitive and physical capacities for independent living and life engagement, is a highly valuable goal” (Judith Gluck et al., *The Gerontologist*, 2025).

## Course Objective

The purpose of this course is to provide an understanding of the concept of healthy aging. Major topics include current biological theories of aging, prevalent diseases and health strategies, Baltimore Longitudinal Study of Aging, psychological factors, social factors, and the nature of healthy aging.

## Accreditation

Continuing Psychology Education Inc. is recognized by the New York State Education Department’s State Board for: Social Work as an approved provider of continuing education for licensed social workers #SW-0387; Mental Health Practitioners as an approved provider of continuing education for licensed mental health counselors #MHC-0080, and licensed marriage and family therapists #MFT-0043; Psychology as an approved provider of continuing education for licensed psychologists #PSY-0006.

## Mission Statement

Continuing Psychology Education Inc. provides the highest quality continuing education designed to fulfill the professional needs and interests of mental health professionals. Resources are offered to improve professional competency, maintain knowledge of the latest advancements, and meet continuing education requirements mandated by the profession.

## Learning Objectives

Upon completion, the participant will be able to:

1. Discuss current biological theories regarding the causes of aging.
2. Acknowledge common older adult diseases and their recommended preventative measures.
3. Articulate findings from the Baltimore Longitudinal Study of Aging.
4. Expound upon psychological effects of aging.
5. Understand social theories of aging, and the value of social support systems.
6. Discuss key characteristics which promote healthy aging.

Director of Operations

Richard Shuman, LMFT

Consultants

Kathryn Wright, LCSW

DeAnne Brining, RN

Dorothy Crawford, Ph.D.

Copyright © 2025 Continuing Psychology Education Inc.

### INTRODUCTION

Aging is a universal human experience interwoven with biological, psychological and social complexities (Guo et al., 2022). In our younger years, we observe illness and death of older family members, with time, we see similar changes in our own generation and ultimately within ourselves. This process is viewed as inevitable and natural, yet, it is also interesting because these changes unfold differently in each of us. Nelson and Dannefer (1992) observed increasing diversity over the adult years on various performance measures suggesting that with age, we become more unlike each other. The principle of *multi-directionality of development* (Baltes & Graf, 1996) suggests that systems within the person develop at different rates - some functions may reveal positive changes and others negative changes over time. Even within the same function, such as intelligence, a person may exhibit gains in one area and losses in another with aging. These individual differences shed optimism and refute the myth that all functioning “goes downhill” as we grow older. In fact, an older adult who exercises and remains active may be stronger, quicker and more mentally agile than a sedentary younger person (Jezewska-Zychowicz & Gajda, 2023; Wickramasinghe et al., 2020).

Though aging is a basic aspect of life, scientific research, including genetic and biochemical bases of disease processes, experimental studies of aging in animals, and evolutionary biology is allowing for greater intervention into the aging process. Such progress is timely as caring for the growing number of elderly is becoming vital. *Life expectancy* - the average number of years of life remaining to a person at a particular age - from birth, has risen in America overall from 62.9 years in 1940 to 76.5 years in 1997, to 75.8 for males and 81.1 for females in 2023 (CDC/National Center for Health Statistics, 2025). From age 65, life expectancy was estimated to be 18 years in 1999 (Hoyert & Murphy, 1999) and 17.5 years for males and 20.2 years for females in 2022 (CDC/National Center for Health Statistics (2024). The number of Americans over age 65 has grown from 3.1 million in the year 1900 (roughly 4% of the population) to over 34 million in 1997 (almost 13% of the population). By 2030, the number is estimated at 69.4 million (about 20% of the population). Further, the oldest-old group, age 85 and older, is expected to rise at the highest rate from currently 1.5% of the population to 2.4% by 2030 to 3.7% by 2040. Not long ago, these advanced ages were not thought possible, yet, interest currently mounts to further increase the human life span and biomedical advances are attempting to delay or eliminate the afflictions of aging.

Factors contributing to longer life span include lower infant mortality, improved prenatal and postnatal medical care, drugs against infectious diseases, and improved nutrition and personal/public hygiene. Research has led to illnesses such as tuberculosis, poliomyelitis, smallpox, measles and many other infectious diseases becoming curable. Despite the convincing increase in longevity, many scientists and medical doctors believe that we die too soon. They are convinced that the

potential life span is about 125 years. Agreed upon is the need to understand the mechanisms of current incurable diseases and of the biological process of aging itself.

Arking (1991) and Moqri (2023) describe aging as cumulative, universal, progressive, intrinsic, and deleterious. Gerontologists, those who study the aging process, agree that aging involves changes in the chemical composition and macroscopic structures of the body; these changes affect the organism’s ability to respond adaptively to the environment - at all levels, from the synthesis of molecules to cognitive abilities (Saxon et al., 2022; Adelman, 1980); and there is greater vulnerability to diseases and to environmental changes and demands leading to a heightened risk of dying (Tenchov et al., 2023; Shock, 1985).

Aging is measured at different levels: the population level examines the life span of individuals; the individual organism level assesses changes in physiological- biochemical functions; the cellular level involves structural and biochemical factors; and the subcellular level focuses on changes in molecules, for example, the activity of enzymatic and repair systems functioning in cells.

Aging is attracting the attention of business and industry, politicians, the media, and the general public due to a rapidly aging society. Data collected by the World Health Organization (WHO) indicates that industrialized countries of the world will be forced to address a phenomenon which humanity has never encountered - demographic changes creating a society dominated in many respects by the needs of an aging or aged population. This shift in the dependency ratio will affect America’s resources and various social structures such as the family, health care, pension and retirement practices, political processes, community and recreational services and housing. As a group, the elderly will have a greater influence on many aspects of society as they spend more money, use more services and possess more political power. An improved understanding of the aging process is recommended to adapt to the changes that will likely occur. To offer greater medical and social opportunities to older people designed to improve aging, in the near future, more physicians, nurses, social workers, counselors, nursing home managers, and many other positions will be needed than are available presently.

### BIOLOGICAL THEORIES OF AGING

Two categories comprise the biological theories of aging: *programmed aging* and *random error*. Programmed aging theories are based on the belief that aging and death are built into the hard-wiring of all organisms. Random error theories believe that aging is due to random damage to an organism over time.

#### PROGRAMMED AGING THEORIES

Genetic life-span theory - It is assumed that “aging genes” exist that count off the years past maturity just as there are “development genes” which lead to the point of maturity in

## HEALTHY AGING

youth. Supportive of this assumption is the fact that life span varies according to species, therefore, life span may be part of an organism's genetic makeup. For instance, a butterfly's life span is 3 months, a human is 125 years, and a giant tortoise 180 years.

Large human population studies in Sweden revealed that longevity has a heritability index (indicates the degree to which a characteristic is inherited) of .26 in men and .23 in women (Herskind et al., 1996a, 1996b). To identify and control the "aging" gene (if it truly exists) could potentially change the nature of aging.

The assumption that length of life is genetically programmed by one or multiple genes controlling the aging process from birth to death, however, is not regarded as viable.

**Genetic postreproduction theory** - This variant of the genetic life-span theory suggests that evolution has selected for species that are vigorous until the time that sexual maturity has passed. Selective survival relates to survival of the fittest when the species can still reproduce, but beyond this time, "good genes gone bad" (Sen, 2025; Hayflick, 1994) take over, in turn, cells are programmed to die.

**Telomere theory** - Proposes that defects develop in gene expression within cells which cause the cells to lose their ability to divide. The loss of ability to reproduce is termed *replicative senescence*. It is known that there are a finite number of times - approximately 50 - that normal human cells proliferate in culture before entering a state in which they cannot further divide (Hayflick & Moorhead, 1961; Zhu et al., 2019; Muraki et al., 2012). Telomere shortening is a fundamental biological process that is closely linked to the aging of cells and organisms, as well as age-related diseases (Vaiserman & Krasnienkov, 2021).

The *telomere* is the terminal region or tail of a chromosome composed of DNA but containing no genetic information; it protects the ends of chromosomes from being degraded and fusing with other chromosome ends, similar to plastic handles that protect the ends of shoelaces. The basis of the telomere theory is that every time a cell replicates, it loses part of its telomere. In other words, older cells (cells that have divided more) have shorter telomeres. This theory believes that cells stop dividing given the short telomere length being sensed as damage to the DNA (Mensa et al., 2019; Herrmann et al., 2018).

Telomere lengths also shorten due to lifestyle factors, stresses and environmental exposures. The National Aeronautics and Space Administration (NASA) observed these factors in the NASA Twin Study which compared telomere lengths in one twin who spent one year in space on the International Space Station (ISS) in contrast to the ground-based twin who is used as a control to isolate spaceflight-related changes. The astronaut-twin experienced a change in telomere length dynamics during spaceflight and within several days of landing. NASA plans on further investigation into environmental effects upon telomere longevity (Edwards

& Abadie, 2019).

Some cells do not display this shortened telomere length and these cells, in particular, contain the enzyme telomerase, which maintains the ends of the chromosomes. Telomerase, it could be argued, can make a cell "immortal," however, some senescent cells continue to manufacture telomerase, hence, this enzyme is not sufficient to prevent cellular senescence (Smith & Pereira-Smith, 1996; Gorbunova et al., 2002).

**Neuroendocrine theory** - Suggests that changes in the hypothalamus, which mediates between the nervous and endocrine systems, are responsible for age changes in the hormones that ultimately lead to aging.

The endocrine glands release hormones into the blood which act on target cells. Hormones regulate many vital activities, including metabolism, reproduction, immune function and growth. In large amounts, hormones are known to be capable of slowing or accelerating some aging processes.

A common precursor to the onset of aging is decline in reproductive capability; the decline in reproductive capacity is governed by the neuroendocrine system, hence, examination of the neuroendocrine system related to aging seems logical.

Research has shown that the longevity of some animals may double when they are raised with a diet which contains all essential nutrients but is low in calories. It is hypothesized that caloric restriction slows aging by retarding the hypothalamic chronometer in the brain. There is evidence to support this hypothesis, however, an explanation of how it works (if it does) is lacking (Weindruch & Walford, 1988).

Another potential neuroendocrine factor affecting aging is a hormone secreted by the adrenal glands called dehydroepiandrosterone (DHEA). This hormone is found in larger amounts in young adults then decreases with age. When administered to mice, DHEA delays immune system dysfunction, increases longevity, and makes the animals "look younger."

Additionally, menopause, which marks the end of the woman's reproductive capacity, exemplifies hormonal events associated with many aging changes throughout the body.

Though the neuroendocrine system affects our bodies in profound ways, there is no direct evidence that it is the cause of all age changes; the exploration of this system's effects on human aging is still at the basic research level (Xing et al., 2023).

**Neuron theory** - Theorizes that loss of neurons regulates the rate of the body's aging. The brain, similar to many other body parts, does shrink in weight and size during normal aging. In fact, brain weight decreases roughly 10%, the convolutions narrow, and the spaces between them widen. The shrinkage may result from loss of neurons, loss of water, or both, however, the cause and relevance of such shrinkage is still debated. In humans, sentiment leans toward the loss of neurons in the brain as the result, not the cause, of aging.

A theory arose suggesting that the brain was the origin of all

## HEALTHY AGING

age changes. The brain was thought to produce a “death hormone.” The hormone is suggested to be made by the pituitary gland and is called the decreasing oxygen consumption hormone (DECO). Donner Denckla, an advocate of the theory, proposed that DECO circulates in the body and disrupts protein synthesis and cell division. These studies as yet are unconfirmed.

Nonetheless, the brain is not being ruled out as the cause of aging. The nervous system does control almost every other system in the body. Further, brain weight is positively correlated with life spans across species. Additional support comes from experiments showing that removal of the pituitary gland, located at the base of the brain, may cause some rejuvenation in animals.

## RANDOM ERROR THEORIES

**Wear and tear theory** - This is one of the earliest aging theories, dating back to 1882, as articulated by the German biologist, August Weismann. He stated, “Death occurs because a worn out tissue cannot forever renew itself.” The theory submits that vital systems accumulate damage from normal use and abuse of daily living culminating in age changes; contributing factors include accidents, disease, radiation, toxins, other detrimental factors, and normal usage. Wear and tear affect the normal biochemical activities of cells, tissues, and organs to the point that our organs and joints simply wear out - such abuse causes aging.

A concern with this theory is that wear and tear is difficult to describe or quantify in most biological systems. We do not know what normal wear and tear is relative to most living cells or the molecules of which they are composed.

Another concern is that most organ systems do not abide by this theory. If organs wear out with continued usage, then athletes should have shorter life spans than sedentary people, but this does not appear to be true (Paffenbarger et al., 1993). In fact, many systems such as the cardiovascular system need regular, moderate exercise to maintain function, and intellectual stimulation may be needed to maintain cognitive functioning (Diamond, 1993). Gerontologists promote the phrase “use it or lose it.”

An exception to this rule is that wear and tear on the joints is a cause of osteoarthritis, and it is very common in late life. Contrarily, total bed rest for the elderly can produce stiffened, immobile joints and problems with many other systems, including cardiovascular, respiratory and neuropsychological systems (Fishburn & de Lateur, 1996). Generally, moderate usage is recommended for organ functioning.

Abuse to a system will shorten its life span as evidenced by skeletal muscular system concerns in professional football players and neurological problems in professional boxers who have sustained many concussions.

Miquel, a biogerontologist, suggests that molecular wear and tear may affect the mitochondria - the “power plants” which provide energy for all the cell’s activities. The mitochondria in old cells of many animal species reveal a decrease in numbers, and an increase in size and various

structural abnormalities. Cultured normal cells reflect these mitochondrial changes at the end of their lifetime as well, prompting Miquel to believe that damage over time to the mitochondria may be a cause of aging.

The major concern with the wear and tear theory as with the mitochondrial version of the theory is the uncertainty whether such damage is the cause or the result of aging. Convincing data exists in favor of and against this theory, but there are many individualized differences even within a single species. While wear and tear occurs, its rate and degree may be affected by lifestyle, genetic makeup, environmental stress, psychological stress, injury, and disease (Sattaur et al., 2025).

**Rate of living theory** - Believes that we are born with a finite amount of some substance, potential energy, or physiological capacity that can be consumed at various rates. Aging begins early in those who spend the energy rapidly and aging is slowed if the energy is exhausted slowly. This theory equates with “live fast, die young.” The theory is traced back to 1908, when Max Rubner, a German physiologist, discovered that various species of animals of different sizes and life spans spent roughly 200 kilocalories per gram of tissue during their lifetime. Larger, long-lived animals spent fewer calories per gram of tissue per year compared to smaller, short-lived species. Raymond Pearl and Ruth DeWitt Pearl, American gerontologists, wrote, “in general, the duration of life varies inversely as the rate of energy expenditure.” Rubner and Pearls proposed that the finite element was metabolic capacity and when it declines age changes occur.

Variants of the rate of living theory suggest that other limiting factors create aging such as the amount of oxygen consumed, number of breaths taken, and number of heartbeats spent. A mouse’s heart, during its three-and-a-half year life and an elephant’s heart, during its seventy-year life, each beat about one billion times; a human heart beats approximately three billion times during its seventy-five year span. Few gerontologists believe, however, that number of heartbeats affects aging.

The rate of living theory lacks empirical backing - the finite substance is unknown and its existence is not certain. Moreover, a centenarian’s heart has beaten about four billion times and humans metabolize roughly 800 kilocalories per gram of tissue in a lifetime - each value about four times that of the oldest elephant - yet, life spans are comparable between the two species. Generally, birds have high metabolic rates and long life spans; human athletes and busy people do not age faster than sedentary people; and among centenarians, some have led active lives, while others inactive. Finally, the literature-supported “use it or lose it” principle defies the notion of not “wasting” yourself by engaging in too much activity.

**Waste product accumulation theory** - Speculates that metabolic waste accumulates in the cells and interferes with normal cell functioning; in time, the toxins and refuse slowly kill the cell. Most waste products are expelled from cells and carried away by the circulatory system and then excreted. In

## HEALTHY AGING

contrast, aging cells accumulate waste matter called *lipofuscin* - a dark mixture of lipoproteins and various waste products which are common in muscle, nerve, and other cells that are not replaced but may remain through the life span. Lipofuscin is so common in old cells that they are called age pigments and are markers for old cells. Lipofuscin production artificially induced in human cells in vitro blocks cell proliferation and induces cell death (von Zglinicki, Nilsson, Docke, & Brunk, 1995). Some speculate that as lipofuscin granules amass in the cytoplasm, they may disturb transport of materials through the cytoplasm or across membranes. They ultimately occupy so much space that there is not enough room for other cellular components.

Despite its presence in some old cells, evidence that age pigments interfere with normal cell function is lacking. The cells do not seem to show signs of distress, even with large amounts of age pigment. Further, some nerve cells and cardiac muscle cells do not contain age pigments, and some old animal cells contain little or no age pigment. Most biogerontologists believe that age pigment is an effect, not a cause of aging (Kim et al., 2022).

**Cross-linking theory** - Postulates that with age, some proteins, including *collagen*, become linked together and may interfere with cellular metabolism by obstructing passage of nutrients and wastes into and out of cells resulting in malfunctioning and aging. Collagen is an important protein found in tendons, ligaments, bone, cartilage, and skin; it makes up nearly 30% of the body's protein.

The process which affects the collagen molecule with age is called "cross-linking." Cross-links are the horizontal strands (similar to rungs of a ladder) that connect the two parallel molecules that form the collagen protein. Over time, rungs of one ladder connect to the rungs of another ladder by forming more cross-links producing larger scaffolds within the tissue. An unknown mechanism prevents this harmful cross-linking in young people but allows its occurrence with age. A manifestation of this process is that our skin becomes more rigid and shrinks in size rather than appearing soft and pliable as in our youth.

This model proposes that cross-links occur within the DNA, creating molecules whose structure is determined by faulty genes. Given accumulation of errors in various molecules over time, age changes result.

Though it is accepted that cross-linking occurs in collagen and some other proteins, it is uncertain whether it exists within the DNA. Further, experimental evidence showing that it affects metabolic processes or causes faulty molecules is lacking. Cross-linking is thought to be one of many biochemical changes contributing to aging but it is not accepted as the main cause.

**Free radical theory** (also called oxidative stress theory) - Believes that a chemical reaction with oxygen forms unstable *free radicals* that unite with susceptible molecules in cells producing age pigments, cross-links in some molecules, and potential damage to DNA. A free radical is an atom or

molecule containing at least one odd or unpaired electron and is chemically reactive because it seeks to bind with other atoms and molecules until obtaining a complete electron pair. The molecule to which the free radical has become attached loses its functioning. Once a free radical is created, it can generate a chain reaction of atoms and molecules stealing electrons from one another. The process of transferring electrons from one molecule to another is termed *oxidation*; generally, the free radical combines with oxygen. The free radical theory suggests that aging is caused by accumulation of irreversible damage due to these oxidizing compounds.

Experimental support revealing the involvement of free radicals in aging comes from *antioxidants* - chemical inhibitors that prevent oxygen from combining with susceptible molecules to create free radicals. Vitamins E and C are considered important antioxidants. Laboratory animals have been fed large amounts of antioxidants and monitored to determine longevity as compared to control animals receiving no antioxidants. Though the results are uneven, research has shown that the antioxidant-fed animals live longer than the controls - increases in longevity up to 30% have been observed in mice. The possibility exists, however, that the antioxidants may have impeded the animals' digestion, resulting in the same effect of calorie-restriction on life expectation. A connection exists between the free radical and caloric restriction theories as suggested by the possibility that the effects of caloric restriction may result from a reduction in the process of free radical formation.

Enzymes called *superoxide dismutase (SOD)* can also destroy some free radicals. Evidence shows longer-living species have higher levels of SOD to detoxify the free radical called superoxide. Humans possess the highest SOD levels of all species studied.

The free radical theory is popular and many gerontologists continue to explore this phenomenon. Additional support is provided by the finding that the rate of production of free radicals is greater in shorter-lived species; perhaps longer-lived species have evolved to produce more antioxidants or free radical destroying enzymes, therefore, reducing the number of damaging free radicals. There is also some evidence that older animals produce more free radicals suggesting a direct link with age. The free radical theory may also be connected to the rate of living and wear and tear theories because all three may be affected by the rate and production of free radicals.

Free radical studies show administering antioxidants to animals seems to postpone cancer, cardiovascular disease, degenerative diseases of the central nervous system, and depression of the immune system. Perhaps antioxidant-fed animals live longer due to postponement of diseases that would have killed them at an earlier age. These same diseases are also postponed by caloric restriction, in turn, skeptics believe that feeding antioxidants mimics calorie restriction. Nonetheless, Sohal and Weindruch (1996) profess that current evidence in support of the free radical theory and future research will support the view that aging results from oxidative stress. Oxidative stress is shown to be related to

## HEALTHY AGING

many pathological states of an organism (Halliwell & Gutteridge, 2015). Oxidative stress is involved in the incidence of chronic diseases involving cardiovascular diseases, cancer, metabolic, renal, lung, and neurological disorders such as Alzheimer's and Parkinson's disease. Cells subjected to various stresses, including oxidative stress undergo cell cycle arrest, termed cellular senescence, and senescence is deemed to be a key hallmark of aging (Jomova et al., 2023).

**Autoimmune theory** - Suggests that the aging immune system loses ability to produce antibodies in adequate number and of the proper type; secondly, it mistakenly produces antibodies against the body's own proteins.

The immune system, one of the body's major defense systems, identifies and destroys foreign substances such as bacteria, viruses, and fungi while leaving normal body cells and molecules undisturbed. Other white cells in the blood produce antibodies which circulate in the blood, deactivate foreign substances (antigens), and prepare them for digestion by other cells.

The *autoimmune theory* states that, with age, the immune system loses ability to distinguish normal from foreign materials leading to the attack and destruction of important body components causing age changes. An example of autoimmune disease in older people is inflammation of the joints (arthritis).

The *immune deficiency theory* believes that the immune system weakens with usage and ultimately cannot defend the body against foreign molecules and microbes. These toxic agents injure and disrupt cell functioning leading to damaging age changes. For example, the immune system of a younger rather than older person may more effectively keep cancer cells in check. A possible cause for immune function decline with age is the thymus gland, located in the upper chest. This gland produces T-cells which are white blood cells that are vital in the body's ongoing effort to fight disease. The thymus begins to wither beyond adolescence and it is theorized that this spurs the ultimate demise of the entire immune system.

The immune system theory has not received strong support for several reasons. First, it is not universal as some animals that age do not manifest a well-developed immune system. Second, the immune system is subject to control by various hormones and the nervous system, therefore, a more basic source could be causing immune function changes with age. Despite the immune system revealing some functional decline with age, it lacks evidence as being the cause of aging.

**Theories of errors and repairs** - Proposes that mutations occur in somatic cells (those not involved in reproduction) causing age changes. A variant of the theory, called the error catastrophe theory (Orgel, 1963), believes that errors in protein manufacturing accumulate exponentially causing many cells to malfunction and die.

The production of proteins and reproduction of DNA are vital to life but these molecules are not always produced with complete constancy. Error theorists submit that an organism's

manufacturing machinery sustains errors, and natural repair processes cannot make perfect repairs every time, and may be flawed from the onset. Errors then affect the molecules that compose or are produced by our cells, metabolic failures arise, and age changes and death result. Evidence shows that errors do occur and repair processes are not perfect and do not function forever. DNA damage affects most, if not all, aspects of the aging phenotype, which potentially makes it a unifying cause of aging (Schumacher et al., 2021).

Preserving the fidelity of DNA is so important that cells have developed methods to repair it given damage. DNA may be damaged by normal background radiation, ultraviolet radiation, cancer-causing chemicals, some normal metabolic processes, cross-linking, and free radicals; fortunately, there are at least six different kinds of DNA repair systems. It is believed that one year's worth of accumulated damage to cells would render cells to be non-functional.

Some research found that DNA repair systems were more efficient in longer-lived species which would support the error hypothesis of aging; perhaps longer-lived species have evolved more efficient ways to correct errors in vital DNA molecules. Confirmation of this finding, however, has not been constant across trials.

Additional research has found evidence that accumulated errors in enzyme proteins may contribute to aging. Speculation is high that some future version of the error theory will explain a number of age changes. It is known that errors do occur in some of the molecules that compose all cells.

**Heat shock proteins theory** - Similar to DNA repair mechanisms which protect genes, *heat shock proteins (Hsps)* represent a type of cellular repair mechanism considered to be an aging decelerator. The name derives from research on effects of heat-related stress upon plants. Hsps are found in every living organism, from bacteria to humans, and in almost all types of cells; they are important in protecting cells from nearly all kinds of stressors, including radiation, infection and oxidation. Additionally, they are important in promoting healthy cell growth and proliferation (Punyiczki & Fesus, 1998).

Jurivich, Qiu, and Welk (1997) suggest that cell aging is defined by poorer responses to physiological stress, possibly mediated by transcription errors in heat shock genes. This reduction in efficiency may interfere with organ functioning and may affect the body's ability to communicate and regulate functioning at the systems level causing failures in homeostasis. Heat shock proteins provide a dual role in cell apoptosis and cell death, thus, further investigation is warranted to explore the role of heat shock proteins in the development of therapeutics for a range of diseases characterized by severe stress conditions and disruption of protein homeostasis (Singh et al., 2024).

**Homeostasis theory** - Organisms must sustain a level of homeostasis, in other words, stability in intracellular and extracellular environmental conditions, such as blood

## HEALTHY AGING

pressure, heart rate, temperature, pH balance, and electrolyte and fluid balance. Conditions change as a function of environmental demands but then must return to baseline levels. Homeostasis demands communication between organ systems and is primarily regulated by the autonomic nervous system via the neuroendocrine system.

Evidence suggests that it becomes harder to maintain homeostasis as we age (Taffet, 2023). Many systems within the elderly reveal slower responses to environmental challenges and more elevated responses requiring much longer to lessen and return to baseline. Both disease and age-related processes are inferred as the cause.

Homeostatic problems may develop from a decline in hormone production or other type of regulatory peptide, target organs may become less responsive, and/or the target organ may synthesize decreased amounts of its product. A number of illnesses associated with aging demonstrate these concerns with homeostasis, such as orthostatic hypotension and diabetes. Research is ongoing.

Order to disorder theory - States that an organism loses efficiency after passing the reproductive age due to a lack of energy needed to maintain the system; increasing disorder causes errors to develop that lead to death.

This model borrows from the field of physics, specifically, thermodynamics. Physicists report that in a *closed system*, matter tends to a state of equilibrium, which equates to increasing disorder. The measure of this state is called entropy and is illustrated in the second law of thermodynamics.

Peak efficiency drops after sexual maturation due to the notion that “perfect order requires infinite work” and a biological system cannot supply infinite work. Further, deterioration and disorder continues to accumulate. Some biogerontologists believe that increased molecular disorder develops from mistakes in molecules (similar to error theories) leading to changes in cells, tissues, and organs called aging.

Humans and animals are examples of an *open system* through which matter and energy flow. We are affected by many environmental factors such as the food we eat and the air we breathe, consequently, it may be argued that the laws of thermodynamics may not generalize well from a closed system to a living organism. Contrarily, our bodies experience the same kind of disorder or entropy in our earlier well-ordered molecules as any machine, complex system, or even the universe itself.

Some biogerontologists argue that aging has many causes requiring a synthesis of many theories while others promote only one theory. Gerontology is still a young science searching for knowledge offering explanation for why we age. Science now knows that age changes occur within individual cells but knowledge has been essentially descriptive - answering “what” happens as we age rather than “why” it happens. Researchers have explained changes that occur as we age from the molecular level to the whole animal but these

descriptive observations do not clarify the basic process. Optimism grows, as many scientists, for the first time in the history of biology, are working in the aging field, and they are encouraged by the potential of modern technological tools, the human genome, stem cell research, and artificial intelligence. Understanding causes of aging may enable scientists to influence the processes of biological aging and to minimize common harmful age changes.

## PREVALENT DISEASES AND HEALTH STRATEGIES

Several major types of physical disease which attack the body’s primary organ systems affect adults. These illnesses are examples of *secondary aging* (also called impaired aging) which constitute later life changes due to disease, as compared to *primary aging* (also called normal aging) which involves age-related changes that are universal, intrinsic, and progressive. People are advised to utilize preventative measures beginning in early adult years to potentially avoid disabling and lethal effects of these conditions.

## CARDIOVASCULAR SYSTEM DISEASES

Diseases affecting the cardiovascular system are the leading causes of death in those over age 65 (Donato et al, 2018), and the leading cause of death worldwide, accounting for 17.7 million deaths per year, and this number is expected to increase to 23.6 million by 2030, which is much higher than cancer and other diseases (Wang et al., 2022). Heart and cerebrovascular disease, together, totaled 43% of all deaths in those over 65 in 1997 (Heidenreich et al., 2011; Hoyert & Murphy, 1999), with over 15 million deaths worldwide (World Health Organization, 1997). Men have a higher chance of dying from heart disease than women; African-American men have the highest death rate from heart disease (more than double the rate of other age and sex groups); African-American women have an elevated death rate for this disease; and Asian/Pacific Islanders have the lowest death rate for heart disease.

Diseases of the heart and arteries are also among the most prevalent chronic condition affecting older adults, specifically, in 1994, over 22 million people in the U.S. reported suffering heart conditions, totaling 30 to 40% of men and women in the over 65 age group.

Presently, four major heart disease risk factors are identified. The first risk factor is leading a sedentary lifestyle, a theme supported by many studies. A large-sample study examined nearly 1400 men from Finland, ages 35 to 63, over an eleven-year interval from 1980 to 1991 (Haapanen, Miilunpalo, Vuori, Oja, & Pasanen, 1996). The 27% of the sample who engaged in vigorous activity at least twice a week revealed a 60% lower death rate from all causes, including cardiovascular disease, compared to those not involved regularly in vigorous activity. Those who burned less than 800 calories per week in some type of physical activity displayed almost five times the risk of dying from

## HEALTHY AGING

cardiovascular disease contrasted to those who burned 2100 calories or more. Further, exercise can improve immune responsiveness in elder adults (Teodoro et al., 2025). Healthy levels of diet and exercise are recommended because a malfunctioning immune system is a common contributor to mortality in middle and later adulthood.

The second risk factor is smoking, and the Finland study typifies common findings. Smoking more than doubled the risk of dying from any disease over the eleven year study and tripled the risk of dying from cardiovascular disease. Researchers do not know the exact cause for smoking increasing heart disease risk but they suspect it damages the arteries and makes them more vulnerable to passing fat cells that contribute to plaque formation.

The third risk factor for cardiovascular disease is body weight (Kannel, D'Agostino, & Cobb, 1996), especially for those between ages 30 and 70 (Stevens et al., 1998). Obesity is defined by the relationship between body weight to body height. The BMI (Body Mass Index) calculates risk based on the weight to height ratio and it equals weight in kilograms divided by height in meters, squared. An ideal BMI is 23 in men and 21 in women while "overweight" is a range of 25 to 29.9 and obesity is 30 and above.

Alcohol intake is the fourth risk factor but findings are inconsistent as to how much is "good" versus "bad." A certain amount (one drink per day) may have some favorable effects on cholesterol levels.

Currently, effective cardiovascular disease treatments include implantable biomaterials and devices, especially minimally invasive interventional ones, such as vascular stents, artificial heart valves, bioprosthetic cardiac occluders, artificial graft cardiac patches, atrial shunts, and injectable hydrogels against heart failure (Wang et al., 2022).

The biomaterials and devices sector has created treatments for various cardiovascular diseases; it is an interdisciplinary field involving medicine, biology, machinery, materials, and other disciplines. Additionally, new clinical technology, advanced clinical surgery techniques, and new materials are projected to develop innovative cardiovascular medical devices. Wang et al. (2022) believe further advanced treatment options will focus on, and offer hope for cardiovascular patients through: 1) biodegradable cardiovascular stents and cardiac occluders with tissue regeneration function; 2) synthetic polymeric valves with biosafety and durability; and 3) injectable hydrogels with tissue regeneration for heart failure therapy.

Additional preventative strategies involve taking dietary supplements, including vitamin C (ascorbic acid), which is found in fruits (Gatto, Hallaen, Brown, & Samman, 1996). Various foods lower LDLs such as high-fiber cereals (Rimm, Ascherio, Giovannucci, Spiegelman, & Stampfer, 1996), and folic acid (Verhoef, Stampfer, Buring, Gaziano, & Allen, 1996), which is found in yeast, liver, green vegetables and certain fruits. Due to heart disease rising significantly in women after menopause, estrogen-replacement therapy is an option. Estrogen seems to lower harmful aspects of LDLs

and increase HDLs (Haines, Chung, Chang, Masarei, & Tomlinson, 1996).

## CANCER

Cancer is a generic term for a group of over 100 different diseases. Each cancer type has its own unique symptoms, treatment, and effect on the individual. From 2010 to 2030, as the American population ages, a 67% increase in the incidence of cancer among Americans over age-65 is projected (Smith et al. 2009). Skin cancer is the most common type of cancer in America. Odds are 40 to 50% of having skin cancer at least once by age 65. Breast cancer is the most prevalent cancer occurring in women and the second leading cause of cancer death (Giaquinto et al., 2024; National Center for Health Statistics, 1997). The most common type of breast cancer is called ductal carcinoma; the cancer starts in the lining of the ducts that lead from the milk-producing glands (lobules) to the nipple. The cancer cells develop in a duct and spread through the wall of the duct and invade the breast fatty tissue. The cancer cells then have the potential to metastasize through the blood and lymphatic system.

Cancer of the reproductive organs most often develops in the uterus, cervix, and ovary in women and in the prostate gland in men.

Many types of cancer become lethal when they spread to the lymph nodes and enter the lymphatic system. Upon reaching these nodes, cancer cells may have spread to other body parts, including other lymph nodes and organs such as the bones, liver, or lungs. Unfortunately, many cancer types have already metastasized by the time the individual becomes aware of his or her condition because many cancers do not cause symptoms when they are growing within the affected organ.

All cancer is genetically caused in that genes that control orderly replication of cells are damaged, frequently due to random mutations that develop in body cells. The mutations arise either as a cell division mistake, in response to injury from environmental agents as radiation or chemicals, or as an inherited tendency for developing certain cancers such as breast and colon cancer. The evolution of a cell from normal to malignant to metastatic appears to follow distinct steps, each controlled by a different gene or set of genes.

Most cancers become more common with increasing age due to age being associated with more cumulative exposure to environmental harmful toxins (carcinogens). The three main cancer risk factors during adulthood are sun-exposure, cigarette smoking, and lack of dietary control. The following cancer statistics were reported in 1998: one million diagnosed skin cancers - many were preventable given use of sun protection; 175,000 deaths caused by tobacco use; 19,000 deaths from excessive alcohol use - often used with tobacco; and 33% of the 564,800 total cancer deaths in 1998 were due to unhealthy nutritional patterns.

Skin cancer is directly linked to exposure to the sun's ultraviolet (UV) radiation. Melanoma is more common in Texas than in Minnesota, where levels of the sun's UV

radiation are weaker. South Africa and Australia have the highest skin cancer rates due to high UV radiation amounts.

Most lung cancer is caused by cigarette smoking and exposure to cigarette smoke is a risk factor for developing cancers of the mouth, throat, esophagus, larynx, bladder, kidney, cervix, pancreas and stomach. Lung cancer risk begins to lessen as soon as one quits smoking. “Second-hand smoke” (exposure to cigarette smoke) is potentially as great or greater a risk for lung cancer.

The link between diet and breast cancer has not been established, but some evidence shows that exercise and a low-fat diet combined with well-balanced meals may be beneficial. Though not clearly established, in men, a diet high in fruits and vegetables may decrease prostate cancer risk while a high-fat diet may increase such risk. Stomach cancer is more common in places where people eat foods preserved by drying, smoking, salting, or pickling, such as Japan, Korea, parts of Eastern Europe and Latin America. Conversely, fresh foods, particularly fruits and vegetables may help to protect against stomach cancer. Colon cancer risk is also considered to be higher in those with a high-fat diet, low fruits and vegetables, and low high-fiber foods such as whole-grain breads and cereals.

Additionally, environmental toxins including pesticides, electromagnetic fields, engine exhausts, and contaminants in water and food are under investigation as possible breast cancer risks. Workplace carcinogens, for example, asbestos and radon (a radioactive gas) increase lung cancer risk; fumes and dust may increase stomach and colorectal cancer. Furthermore, one’s lifestyle, history of disease, variations due to race and ethnicity, and hormonal factors may increase cancer risk.

The American Cancer Society recommends frequent cancer detection screenings such as breast self-examination and mammograms for women, prostate examinations for men, and colon cancer screenings for men and women.

Cancer treatment modalities include: surgery, radiation therapy (high-energy X-rays which damage cancer cells and stop their growth), chemotherapy (use of drugs to kill cancer cells), biological therapy (involves substances called biological response modifiers that assist the body’s immune system in fighting disease), allogeneic hematopoietic stem cell transplantation, pharmacological hormone therapy, treatments targeting genes with oncogenic alterations and related signaling pathways, photodynamic therapy, antibody drug conjugates, immune check point inhibitors, bispecific T-cell engagers, oncolytic virus therapy, and chimeric antigen receptor T cell therapy (Sonkin et al., 2024). New treatment methods are anticipated over the next few decades as cancer research evolves, and cancer deaths may decrease given educational programs targeting high-risk groups on preventable cancers such as lung cancer. Future cancer treatment options may include: cancer vaccines, microRNA, short hairpin RNA, and short interfering RNA based cancer therapeutics, pulsed electromagnetic field therapy, preferential homing of some stem/progenitor cells and some bacteria to tumors, and tumor innervation. A number of gene therapy

approaches include: ribozyme-based RNA reprogramming, using CRISPR to target tumor-specific repeat sequences and tumor-specific junctions created by somatic structural alterations, using somatic mutations as targets for insertion of sequences coding for peptides known to trigger immune response due to previous routine vaccination such as measles or mumps, coding for peptides triggering cell death, delivery of nanoparticles to different organs, and improvements in the efficiency of gene editing (Sonkin et al., 2024).

## MUSCULOSKELETAL SYSTEM DISORDERS

The two main musculoskeletal system disorders affecting middle-aged and older adults are arthritis and osteoporosis (Grote et al., 2019). The effects of these common ailments range from minor but annoying limitations to severe disability.

Arthritis is a general term applying to conditions which affect the joints and surrounding tissues. The most common form of arthritis is termed *osteoarthritis*, and it is one of the most frequent chronic conditions in older adults, affecting about 50% of those over age 65 and more women than men. This condition develops in joints that are injured due to overuse or from being overweight. Ultimately, this injury or repeated impact thins or wears away cartilage designed to cushion the ends of the bones in the joint leading to the bones rubbing together. The synovial fluid which fills the joint loses its shock-absorbing properties and joint flexibility is lowered, bony spurs develop, and the joint swells. These alterations in joint structures and tissues cause pain and loss of movement.

Pain medications are often prescribed, including aspirin, acetaminophen, ibuprofen, and aspirin-like drugs termed nonsteroidal anti-inflammatory drugs (NSAIDs). The injection of corticosteroids are sparingly used to reduce swelling and inflammation. While these pain medications only lessen symptoms, new treatment techniques are becoming available such as injection of a synthetic material into the arthritic joint designed to replace the loss of synovial fluid. Another technique involves injection of sodium hyaluronate into the joint which is an injectable form of a chemical usually present in high amounts in joints and fluids. Total replacement of an afflicted joint such as a hip or knee is also possible.

*Rheumatoid arthritis*, an inflammatory disease causing pain, swelling, stiffness, and loss of function in joints, is an *autoimmune disease* whereby the immune system destroys the person’s own cells inside the joint capsule. The wrist joints and finger joints closest to the hand are often affected. Activity that induces stress on the joints does not appear to affect this condition as is the case with osteoarthritis. Whereas osteoarthritis is a progressive degenerative disease, rheumatoid arthritis varies in course and may last from several months to one or two years but can persist for many years in some people. Life expectancy of those with rheumatoid arthritis is reduced by seven years in men and three years in women.

Roughly two million people in America and 165 million

## HEALTHY AGING

people worldwide have rheumatoid arthritis; this equates to 460 per 100,000 population from 1980 to 2018 (Almutairi et al., 2021; World Health Organization, 1997), and it is two to three times more likely in women. The cause is a combination of genetic and environmental factors and it can be triggered by exposure to an infectious agent.

Treatment focuses upon relieving pain, lessening inflammation, and slowing or stopping further joint damage. Drugs may include disease-modifying antirheumatic drugs (DMARDs), additionally, corticosteroids, aspirin or NSAIDs may be recommended, and joint replacement or tendon reconstruction may be an option.

Normal aging associates with loss of bone mineral content due to imbalance between bone resorption and bone growth. Age-related changes in body composition, such as loss of bone mineral content (Boonen et al., 1996), fat increase, and muscle mass decrease (Bjornstorp, 1996) are considered endocrine system related. *Osteoporosis* (means “porous bone”) is loss of bone mineral content greater than 2.5 standard deviations below the mean of young white, non-Hispanic women.

Among American individuals 50 years of age and older, 10.2 million have osteoporosis, additionally, 43.4 million have low bone mass (Wright, et al., 2014). There has been a statistically significant increase in osteoporosis among Americans 50 years of age and older from 2005 to 2018 (Naso et al., 2025). Osteoporosis occurs in 4% of women aged 50 to 59, in 19% in their 60s, 31% in the 70s, and in 50% of women over 80 (National Center for Health Statistics, 1997). Women experience greater risk than men because they have lower bone mass and menopause accelerates the process due to its estrogen production decrease.

The risk of developing osteoporosis is increased by alcohol and cigarette smoking and is decreased by adequate intake of calcium through dairy products, dark green leafy vegetables, tofu, salmon, and calcium-fortified foods such as orange juice, bread, and cereal. Vitamin D, as a dietary supplement or by exposure to sunlight is a preventative source because it is involved in calcium absorption and bone health. Exercise and physical activity also reduce osteoporosis risk.

Osteoporosis treatment attempts to restore bone strength through nutritional supplements and a weight-bearing exercise program. Medication can be used to slow or stop bone loss, increase bone density, and lower fracture risk. Hormone replacement therapy has been a successful drug in osteoporosis treatment.

## DIABETES

This disease is caused by a defect in the metabolizing of *glucose*, a simple sugar which is a vital source of energy for the body’s cells. In the most common type of diabetes, Type 2, or noninsulin dependent diabetes (NIDDM), the pancreas produces some insulin, however, the body’s tissues do not respond to the insulin signal, a condition termed insulin resistance. Insulin does not bind to the cell’s insulin receptor,

and glucose is not transported into the body’s cells for use. The excess glucose overflows into the urine and leaves the body causing loss of a main energy source.

Symptoms include fatigue, frequent urination, thirst, weight loss, blurred vision, frequent infections, and slow healing of sores. Hypoglycemia occurs when blood sugar levels become too low causing one to potentially feel nervous, jittery, faint, and confused. Given this condition, the individual must eat or drink something with sugar in it as soon as possible. Hyperglycemia occurs when blood sugar levels rise too high and can also cause serious illness.

There are approximately ten million Americans diagnosed with diabetes and possibly five million more who have the disease but have not been diagnosed. In 1997, over 62,000 people in America died from diabetes and 76% of these deaths were in the over 65 age group (Hoyert & Murphy, 1999); over 95,000 Americans died from diabetes in 2023 (Ahmad, 2024). The incidence is 60% higher in African Americans and about 115% higher in Mexican Americans and Puerto Ricans as compared to whites.

Management of Type 2 diabetes includes diet control, exercise, and frequent blood testing to monitor glucose levels; the goal is to keep blood sugar within acceptable levels. Taking oral drugs or insulin to lower blood glucose levels may be needed. An exercise program designed to manage weight and lower blood pressure and blood fats can produce reductions in blood sugar levels.

## ALZHEIMER’S DISEASE

Alois Alzheimer, a German neurologist, was the first person to correlate brain tissue changes with observable dementia symptoms. Alzheimer treated a patient, Auguste D., who experienced progressive mental deterioration, increasing confusion and memory loss. Through a staining technique, he observed an odd disorganization of the nerve cells in the patient’s cerebral cortex and, in 1907, he published in a medical journal that these microscopic changes caused the dementia. Recent analysis of this patient’s brain slides confirm the correlation (Enserink, 1998). A definite diagnosis of Alzheimer’s disease (AD), for almost a century, has always required an autopsy revealing these characteristic neurofibrillary tangles (tangled nerve fibers) and beta-amyloid plaques (looks like accumulated waste products of collections of dead neurons), however, in 2007, an international working group (IWG) of dementia experts described biomarkers as evidence for the diagnosis of AD. Biomarkers for AD is defined as “signs of the physical changes in the brain that contribute to disease progression.” In 2018, a group sponsored by the Alzheimer’s Association and the U.S. National Institute on Aging reported that biomarkers did not only confirm a diagnosis of AD but biomarkers were a diagnosis of AD. This process characterizes AD as a condition displaying abnormal protein deposits in the brain that starts with an asymptomatic phase and progresses to mild cognitive impairment (MCI) and then to a level of dementia that interferes with the processes of daily living. Researchers

are trying to identify AD biomarkers and targets, and in the process, they are utilizing machine learning and deep learning computational methods (Alamro et al., 2023).

AD, a progressive neurodegenerative condition, affects approximately 6.7 million Americans aged 65 or older in the year 2023. An estimated 5 to 7 million Americans aged 65 and older may have mild cognitive impairment (MCI) due to AD (the MCI will likely advance to Alzheimer's disease dementia). By year 2050, those with AD is projected to double in the U.S. and Europe, and triple worldwide (Monfared, 2024).

Psychological symptoms evolve gradually over time beginning with early signs of occasional memory loss for recent events and familiar tasks (Knopman et al., 2021). Cognitive functioning changes represent the essence of the disease but personality and behavior changes ultimately arise. The final stage involves inability to perform simple and basic everyday functions.

AD is known to be associated with the formation of plaques and tangles, especially in brain areas controlling memory and other important cognitive functions. The cause is unknown, but a common theory is that genetic abnormalities are responsible for neuron death (Wisniewski, Wegiel, & Kotula, 1996). New genetic engineering technology is discovering genes that may cause the brain changes associated with this disease. For example, the *apolipoprotein E (ApoE) gene*, located on chromosome 19, may be involved in plaque formation.

Genetic contributions to AD are thought to explain about 40 to 50% of the disease, in turn, environmental factors are studied. Serious injuries involving loss of consciousness increase the risk of acquiring the disease in late life. Such injury may initiate production of cytokines and other proteins, a process which ultimately destroys neurons (Griffin et al., 1998).

Higher education and continued mental activity throughout life may protect against AD. A study of 678 nuns within a religious order known for low rates of AD revealed that they lived an intellectually challenging life into their 80s and 90s. One nun maintained high scores on cognitive tests until her death at age 101. Amazingly, this person's autopsy showed characteristic plaques and tangles found in the brains of those with profound behavioral deficits (Snowdon, 1997). The study also highlights that some people can function through life without any observable signs of the disease.

At present, there is no cure for AD. The drugs currently being tested have not offered significant improvements in Alzheimer's patients, but they may lead to more productive treatments. AD is complex due to numerous genetic and environmental risk factors, and different clinical presentations which suggests it is more similar to a syndrome than to a traditional disease, given its pathological manifestation displaying a convergence of pathogenic pathways. As such, a solution needs a multifaceted approach rather than a 'silver bullet' - coordinating multiple strategies simultaneously may be required (Korczy, 2024).

## BALTIMORE LONGITUDINAL STUDY OF AGING

The Baltimore Longitudinal Study of Aging (BLSA) began in 1958, and is a longitudinal study conducted by roughly 150 gerontologists examining physical, mental, and emotional aging effects in healthy people. Nathan Shock, often regarded as the father of gerontology in the United States, is accredited with initiating the study. Participants include over 1500 men and 700 women, ranging from ages 17 to 96, who undergo extensive testing over two and a half days every two years. Subjects are mostly healthy, well-educated and working in or retired from high-level positions and have been followed for an average of 13 years. The following is a review of BLSA findings regarding aging changes.

Appearance - Both sexes lose about one-sixteenth of an inch per year in height starting at about age 30.

With age, the extremities become thinner and the trunk thicker.

Dental - The chance and severity of periodontal disease increases with age.

Weight and metabolic changes - Life expectation is not greatest for the leanest, rather, for those ranging from the middle of the "desirable" weight range to 20% over that midpoint.

Weight lowers between age 55 and 75 due mainly to loss of lean tissue, muscle mass, water, and bone.

Blood cholesterol rises from early adulthood to age 60 or 65 and then falls.

Coronary artery disease risk factors are cigarette smoking, hypertension, elevated serum cholesterol and low density lipoproteins, low vital capacity, diabetes, and obesity. The incidence of this disease increases with age.

If disease-free, an older person's heart pumps about as well as that of a young adult. Evidence does not show that heart function declines with age.

Pulse rate diminishes with age and is not a health problem.

Reaction time - Those over age 70 show a declined ability to detect and report small environmental changes.

With age, responses to stimuli become slower and have a greater chance of being inaccurate, particularly complex tasks.

There is a 20% slowing in reaction time from age 20 to 60.

Cognitive - Short-term memory declines as we age.

Beyond age 70, performance on logic tests decreased for most BLSA participants, but no change occurred for some.

Ability to learn oral material decreases only in those over 70.

Those who experience no decline on mental tasks exist in every age group, including the oldest.

Visual memory, measured by reproducing geometric designs from memory, slightly declines between ages 50 and 60, and diminishes greatly after 70.

Vocabulary scores show no change with age.

Personality - Barring disease, personality traits remain basically the same throughout life. The notion of becoming crankier or more mellow with age is a myth. Preference for fast-paced activities decreases around age 50. Evidence suggests that older people do not become hypochondriacs.

## HEALTHY AGING

Sexuality - Sexual activity decreases with age despite maintenance of normal amounts of sex hormones. A slightly higher testosterone level is present in more sexually active men.

Sperm counts per unit volume or per ejaculate remain the same, but the proportion of immature sperm present rises with age.

Prostate gland enlargement is common in men over age 60.

Sexual daydreams lower in frequency and intensity with age and they virtually disappear after age 65.

Relative frequency of sexual activity does not change with age; the most sexually active men in their 70s were also highly sexually active in their 20s.

Senses - Visual acuity decreases with age, but wearing glasses allows for 20/20 vision or better into the 80s.

There is a progressive loss in ability to hear sound at all frequencies.

With age, ability to taste sweet or sour does not change, but it becomes a little more difficult to detect salty or bitter.

Ability to identify odors declines with age.

Physiological - Kidney function, measured by ability to clear nitrogenous wastes from the blood, decreases with age.

Lymphocytes, a white blood cell, show a lowered ability to kill cancer cells starting at age 40. Other white blood cells which fight infectious disease, termed neutrophils, also decline in ability with advancing age.

Pulmonary function declines with age.

Physical activity and maximum exercise performance decline with age. Physical performance can improve over time given lifestyle changes such as daily exercise. Physical performance can delay, eliminate, or slow some diseases, but evidence does not reveal that increased exercise can affect basic aging causes.

Gender aging differences - Pre-menopausal women reveal no bone density loss with age, but post-menopausal women have a faster rate of bone loss than men of similar age.

Aging men and women have different proportions of the several types of white cells found in the bloodstream.

Older men have 20% higher maximum oxygen consumption capacity than women.

The longitudinal decrease in hearing sensitivity is more than twice as fast in men than women.

The BLSA offers the following conclusions:

a) Chronological age alone is a poor predictor of performance because age changes are individualized - an 80 year-old may perform equally well as a 50 year-old; b) Some decrements are inevitable effects of aging, such as reduced reaction speed and short-term memory losses; c) Many disabilities associated with old age may be caused more by effects of disease than aging; d) Aging results from interaction of genetic, environmental, and lifestyle factors; e) Lifestyle decisions, such as maintaining a low-cholesterol diet or stopping cigarette smoking, can affect occurrence or progression of some age-related diseases.

## PSYCHOLOGICAL FACTORS

The effects of aging upon mental abilities may influence ability to adapt to everyday life. Further, cognitive abilities can affect self-esteem and how we view our own aging process. Despite some losses in speed and memory in later adult years, normal age-related changes are not completely negative. Fortunately, individuals can compensate for memory changes by implementing a balanced approach to their self-concept.

Information Processing - Age-related changes in information processing are determined by measuring *psychomotor speed* - time needed to process a signal, prepare a response, and execute the response. Research shows that reaction time increases with age in adulthood. Normal healthy aging leads to various sensory and cognitive changes, including loss of sensory acuity, and reduced processing speed (Jones & Noppeney, 2024). The *general slowing hypothesis* states that the increase in reaction time is due to a general decline of information processing speed within the aging nervous system. The *age-complexity hypothesis* suggests that due to slowing of central processes in the nervous system, age differences increase given increasing task complexity (Cerella, Poon, & Williams, 1980). Older adults perform at comparable speeds on tasks which can be completed quickly by a young adult (500 milliseconds), but on tasks requiring 1000 milliseconds for young adults, older adults take proportionally longer (1500-2000 milliseconds). The general slowing hypothesis indicates that older adults do not become deficient during information processing tasks, rather, they are just slower.

Attention - Healthy aging often involves impaired attentional processes (Jones & Noppeney, 2024). The slowing of reaction time with age may have many causal factors, including the possibility that older adults have difficulty during information processing when the stimulus to be focused upon is placed into the system. *Attention* involves ability to focus or concentrate on a portion of experience while ignoring other features of experience, shifting that focus as required by the situation, and coordinating information from multiple sources. Additional cognitive operations, such as memory or problem-solving, may then be performed on this information.

The theory of *attentional resources* and aging suggests that older adults have limited available energy for cognitive operations due to reductions in central nervous system capacity (Salthouse, 1985). The *inhibitory deficit hypothesis* proposes that inhibition (ability to restrict attention to irrelevant or distractor information) is negatively affected by aging, while activation (ability to engage the search process) is spared aging effects. Studies reveal that younger adults inhibit response to distractor information better than older adults indicating that older people are less likely to inhibit processing of irrelevant information. These theories imply that older adults focus more effectively given distractions kept

to a minimum.

Memory - This construct is simplistically defined as the acquisition, storage, and retrieval of information. It involves a sequence of stages allowing information to be initially processed, maintained in a holding pattern, then either discarded or moved into more or less permanent storage. Most researchers agree that memories are based on patterns of neuronal activity in the brain.

Previously called short-term or immediate memory, *working memory* is a system which keeps information temporarily available and active while the information is being used in other cognitive tasks (Baddeley, 1986). Three components are theorized to be associated with working memory. The *phonological loop* is composed of a memory store for speech-based information called the phonological store, and an articulatory control process for processing this information. For example, one may hear a name (phonological store) and repeat it several times subvocally (articulatory control process) to avoid forgetting the name. The *visuospatial scratch pad* facilitates manipulation and maintenance of visual and spatial images, for instance, it may be used to determine the shortest route between home and a grocery store. The third component, the *central executive*, determines how to allocate cognitive resources such as whether to rehearse that name or to create a mental street map in deciding the shortest route.

Most studies on working memory span reveal clear age-related deficits, both in verbal (Bugg et al., 2007; Verhaeghen, Marcoen, & Goossens, 1993) and computational span (Salthouse & Babcock, 1991). It appears that the articulation rate slows in adulthood which may affect processing of information in the phonological loop (Multhaup, Balota, & Cowan, 1996). The visuospatial scratch pad diminishes with age as apparent from spatial memory deficits in older adults (Tsvetanov et al., 2013; Smith, 1996; Wilkniss, Jones, Korol, Gold, & Manning, 1997).

Information does not always remain within working memory, eventually, it is either forgotten or consolidated in *long-term memory* - the repository of information held for a time-frame ranging from several minutes to a lifetime. The effect of working memory deficits on long-term memory is illustrated in the *environmental support hypothesis* which proposes that age-related differences exist on tasks that provide little context or support and require high levels of self-initiated processing in which the person must work hard to remember the material ( Craik, 1994). When such high demands are placed on working memory, older adults do not process the material as efficiently or effectively.

Memory for information which is highly familiar and frequently retrieved, and memory that occurs without conscious processing are immune to negative aging effects. The following types of memory, therefore, show little or no age effects: semantic memory - the equivalent of "knowledge," and includes the words and definitions of words in one's vocabulary or storehouse of historical facts; long-term semantic memory ability actually increases with

advancing age relative to the performance of 25- to 29-year-old individuals, with higher mean scores for age cohorts through the 75-79 age group (Larrabee, 2019; Salthouse, 2010); procedural memory - non-verbal memories in the form of knowledge of how to perform certain activities such as riding a bicycle; implicit memory - recall of information acquired unintentionally; autobiographical memory - recall of information from one's own past (no aging effect if the information is of great importance); and prospective memory - recall of events to be performed in the future (no aging effect for event-based prospective memory such as remembering to meet someone after dinner, but there are significant age effects in time-based prospective memory such as remembering to meet someone at the specified time of 6:00 p.m.).

Along with working memory span, the following types of memory have significant age effects: episodic memory - memory for events (episodes) and can include recall of information administered during a memory experiment such as a word list; a progressive decline in long-term verbal and visual episodic memory begins in the age 45-54 cohort (Larrabee, 2019); flashbulb memory - remembering details of a distinctive historical event; source memory - recall of where information was heard or seen; tip-of-the-tongue phenomenon - difficulty in retrieving a well-known piece of information such as a name or word; and remote memory - recall of information from the distant past. It appears that memory tasks that require the type of processing involved in working memory, which put high demands on cognitive resources, are negatively affected by aging. Evidence from brain imaging and other neurophysiological data suggest possible areas of age-related effects are the frontal lobes, which may affect working memory and source memory (Trott, Friedman, Ritter, & Fabiani, 1997), and the hippocampus-medial temporal lobe areas (Henkel et al., 1998; Raz, Gunning-Dixon, Head, Dupuis, & Acker, 1998; Smith, 1996).

Older people participating in high levels of aerobic fitness reveal a small (5%) but significant improvement on complex speed-based cognitive measures (van Boxtel et al., 1997). An older adult Swiss sample study found that high levels of antioxidants in the blood (ascorbic acid and beta-carotene) correlated with enhanced memory functioning over a 22-year period (Perrig, Perrig, & Stahelin, 1997), but this finding requires additional confirmation.

A healthy lifestyle is correlated with slower memory decline, even in the presence of the apolipoprotein E epsilon 4 (APOE4) allele, which is a genetic risk factor for Alzheimer's disease (Jianping et al., 2023). Simple practice has been shown to improve memory task performance and negate negative effects of mental inactivity (Lachman, Weaver, Bandura, Elliott, & Lewkowicz, 1992). Training older adults in traditional memory improvement techniques such as mnemonics has offered limited success, instead, helping them to devise their own memory enhancement strategies has yielded better results (Park, Smith, & Cavanaugh, 1990). Self-guided practice offers a more positive training effect (Kotler-Cope & Camp, 1990; Willis, 1990), and improves the

older person's feelings of efficacy and mastery (Cavanaugh & Green, 1990).

**Language** - The average healthy older adult does not experience significant losses in language usage (Burke, 1997). Age-related changes in cognition, however, may affect language usage (MacKay & Abrams, 1996). Working memory declines create difficulty in gaining meaning from written material or conversation because information must be retained in memory while new information is being processed. Cognitive processing declines are viewed as a significant factor in reducing the quality of interpretations which older adults derive from written and spoken language. It may be more difficult to retrieve needed words to complete thoughts or sentences while speaking (Burke & Shafto, 2004; Shafto & Tyler, 2014; Kemper, 1992), and an older adult's speech may become more ambiguous (Shadden, 1997). Additional difficulties include word identification in speech (Thornton & Light, 2006), use of specific (not vague) vocabulary (Burke & Shafto, 2004), production of propositionally and syntactically complex speech (Shafto & Tyler, 2014) with complex idea density (Peelle, 2019), and maintenance of targeted verbosity, specifically, topic-centered discourse (Thornton & Light, 2006).

Older individuals can compensate for aging effects on speech production and understanding through their backlog of experiences. They can utilize the context of a situation to understand the correct meaning, and they can anticipate and organize information, at times, better than a novice in that situation. Compensatory strategies both support declining memory features and ensure preserved functions. Neurocognitive compensatory strategies are essential for language (Park & Reuter-Lorenz, 2009). Compensatory strategies recruit additional neural resources as a relevant adaptation that promotes accuracy in a task (Peele, 2019).

Two well-known models attempt to explain how compensatory strategies support aging cognition and language. First, the compensation-related utilization of neural circuits hypothesis model (CRUNCH) indicates that older individuals recruit additional neural resources to cope with increasing cognitive load and demand (Reuter-Lorenz & Lustig, 2005; Reuter-Lorenz & Cappell, 2008). Second, the scaffolding theory of aging and cognition revised (STAC-r) model believes that elders develop complementary neural circuits to achieve cognitive goals, specifically, compensatory abilities develop over the entire life course (i.e., in a scaffolding way), and these strategies are more prominent during the later life stage (Park & Reuter-Lorenz, 2009; Reuter-Lorenz & Park, 2014).

Aging speakers are able to preserve the formal aspects of language, such as syntactic form (Wingfield & Grossman (2006), and benefit from semantic priming and contextual cues (Diaz et al., 2016).

**Problem-solving** - The process of problem-solving involves assessing the present state of a situation, determining the desired end-state, and finding ways of changing the current

into the desired state. Through years of problem-solving experience, an older adult may routinely search for relevant factors in a problem, and this increased selectivity to information can reduce the risk of becoming burdened with excess information. Secondly, an experienced problem-solver often has well-organized storehouses of knowledge that can be easily accessed and implemented. Supportively, Chen et al. (2017) and Strough et al. (2018) found that older adults increasingly utilize knowledge, skills obtained through life experiences, education, and learning (crystal intelligence) to assist in everyday problem solving, while young adults rely almost entirely on the ability to reason, solve novel problems, and adapt to new situations, thus, processing and applying new information flexibly (fluid intelligence).

Research on speed of decision-making shows that older adults can arrive at an answer more quickly than younger persons who do not have their knowledge bases as well-stored or categorized. It also appears that older adults make quicker decisions in areas that they are not especially expert; they are also less likely to seek additional information upon making the decision compared to young adults (Meyer, Russo, & Talbot, 1995). Quicker problem-solving in older adults may result from their greater experience base creating a feeling of lower dependency on incoming information compared to younger adults.

Two problem-solving styles are the *top-down approach* - the person uses "heuristics" or rules-of-thumb to approach a problem (this method is commonly used by an expert problem-solver as illustrated by a chef who does not need to measure the sugar needed in pie crust), and the *bottom-up approach* - the individual collects as much data as possible before making a decision (the novice baker would precisely measure the needed sugar for pie crust). The top-down approach is quicker but it can lead to mistakes given incorrect assumptions during the beginning stage of problem-solving. This type of mistake occurs given too much, not too little familiarity. Young adults use more bottom-up processing, older adults use more top-down, and middle-aged adults use the better approach of combining both (Sinnott, 1989).

Willis (1996) theorizes that older adults rely on top-down processing and make quicker decisions with less information as an attempt to conserve cognitive resources. Further, this style may lessen the elder's discomfort of ambiguity and the quicker decision may give the individual more time to prepare for necessary action given their decision.

Due to greater experience and expertise, older persons have an advantage given familiar choices, but they are disadvantaged when a familiar concern arises with a change, or when a premature decision results in ignoring important information. Further, decisions by older adults may be negatively impacted when the cognitive demands of decisions exceed deliberation capacity given age-related declines in fluid cognitive ability (Del Missier et al., 2012, 2013). Decisions by older adults may benefit from cumulative life experience (Eberhardt et al., 2018; Li et al., 2013, 2015), however, knowledge deficits in specific areas can produce negative consequences (Stewart et al., 2018). In contrast,

elders are adaptive decision makers (Mata et al., 2012) as they actively lower the complexity of information via their search strategies (Besedes et al., 2012), and selectively utilize cognitive resources for personally important decisions (Hess & Queen, 2014). The enhanced emotion regulation by older adults helps to compensate for cognitive declines (Bruine et al., 2014; Eberhardt et al., 2018; Mikels et al., 2010). More research is needed to strengthen the aging decision makers' capacities and create interventions that facilitate daily life decisions (Strough, 2020). Young problem-solvers may make mistakes due to unfamiliarity with many situations, however, their ability to process larger amounts of information in a shorter time can lead to avoiding some mistakes that older adults may make.

Intelligence - The construct of *intelligence* has been difficult to define beyond the idea that it represents the overall quality of the individual's mental abilities. Results from the Wechsler Adult Intelligence Scale, developed by David Wechsler in the 1930s, indicated that age-related differences in intelligence followed the "classic aging pattern" (Botwinick, 1977) of an inverted U-shaped pattern; specifically, a peak in early adulthood followed by steady decline over succeeding decades of adulthood. This conclusion has been challenged by the *Seattle Longitudinal Study (SLS)* in association with K. Warner Schaie which found, for most abilities, there was an increase or no change between the first and second testing, even within the oldest age group. The archives of the SLS are considered the major repository of data on adulthood intelligence.

Breit et al. (2024) assessed the stability of cognitive abilities, including general intelligence, and found support for the SLS determination that intelligence is stable over time. Cognitive abilities, including general intelligence, along with domain-specific abilities as fluid reasoning, comprehension knowledge, working memory capacity, and processing speed, are considered to be classified as some of the most stable psychological traits. Limited research exists, however, regarding the stability changes of these factors over age and time interval, and across abilities, tests, and populations. Determining the conditions under which cognitive abilities vary in stability is relevant to theory development and applied contexts where cognitive assessments affect decision making about treatment and interventions. Breit et al. (2024) presented a meta-analysis using data from 205 longitudinal studies that included a total of 87,408 participants, resulting in 1,288 test-retest correlation coefficients among the relevant variables. Results indicated low stability in cognitive abilities in preschool children, rapid increases in stability in childhood, and consistently high stability from late adolescence to late adulthood. This same pattern described age trends in stability after adjusting for test reliability. The most stable cognitive ability was found to be general intelligence, but many specific cognitive abilities are likewise stable.

Multi-dimensional intelligence theories suggest that intelligence involves multiple abilities or dimensions of abilities rather than the idea of intelligence as a unitary

construct. The multi-dimensional approach proven to be effective in understanding intelligence is the primary mental abilities framework presented by Thurstone (1938). This model proposes seven *primary mental abilities*, including verbal meaning, word fluency (generating words following a particular lexical rule), number (arithmetic), spatial relations, memory, perceptual speed, and general reasoning. Five of these seven abilities form the basis of most studies on adult intelligence: Verbal Meaning, Space, Reasoning, Number, and Word Fluency.

Stemming from the primary mental ability theory, Raymond Cattell (1963) proposed that intelligence involves two basic sets of abilities called *secondary mental abilities*: one set is based on unlearned thought processes and the other set on educational training. The first ability, *fluid intelligence*, represents the individual's innate abilities to perform higher-level cognitive operations involving integration, analysis, and synthesis of new information, "the sheer perception of complex relations" (Cattell, 1971, p. 98). Fluid intelligence also involves the quality of biopsychosocial factors such as nervous system functioning and sensory structures and it cannot be trained or taught, rather, it is a "pure" measure of ability not influenced by educational experiences. The second ability, *crystallized intelligence*, comprises acquisition of specific skills and information acquired by familiarity with the language, knowledge, and conventions of one's culture; it involves the learned ability to infer relationships, make judgments, analyze problems, and utilize problem-solving strategies. Together, fluid and crystallized intelligence include biological factors related to the nervous system, psychological factors encompassed in cognitive processing, and social factors obtained from education and experience in one's culture.

The peak of fluid intelligence is theorized to occur during the years of adolescence, when theoretically, the nervous system and sensory structures are at optimum levels. Beyond adolescence, age-changes which diminish these systems create a downward trajectory in fluid intelligence. Crystallized intelligence continues to grow throughout adulthood while the person gains experience and culture-specific knowledge.

Verbal meaning scores peak by the 50s on cross-sectional studies and somewhat later on longitudinal studies, then scores start to drop about 10 years later followed by performance being about half the peak-level by age 80. Numerical ability peaks in middle age and undergoes a detectable drop by the 60s. The three primary mental abilities representing fluid intelligence, spatial orientation, inductive reasoning, and word fluency remain stable until the late 60s and drop steadily afterward.

The SLS reveals that health status affects intelligence test performance. Arthritis, cancer, and osteoporosis are associated with lower intelligence test scores (Schaie, 1996), as is cardiovascular disease, and hypertension in middle adulthood is a risk factor for poorer cognitive performance in the 70s and beyond (Launer, Masaki, Petrovitch, Foley & Havlik, 1995).

Gender differences reveal that men outperform women on

numerical skill, the crystallized ability of knowledge of general information, and the fluid ability scale of spatial orientation. Women score higher on a fluid measure called Digit Symbol - substituting symbols for digits in a speeded coding task (Kaufman, Kaufman, McLean, & Reynolds, 1991; Portin, Saarijaervi, Joukamaa, & Salokangas, 1995). Changes through adulthood indicate that men show earlier losses on crystallized abilities and women decline earlier on fluid abilities (Dixon & Hulstsch, 1999).

SLS data reveal that those with higher levels of education are protected somewhat from aging effects on intelligence, and this buffering effect also includes being involved in a complex and stimulating work environment, being married to a spouse with higher levels of education and intelligence, and exposure to intellectually stimulating environments in general. Retirement positively affects maintenance of intellectual functioning if one is leaving a boring and routine job, but those leaving a complex and stimulating occupation sustain a more pronounced decrement after retirement (Schaie, 1996). Cumulatively, these qualities produce higher amounts of "Life Complexity" (Schaie, 1983).

Older adults who demonstrate more flexible attitudes and personality style are less likely to experience a decline in intellectual functioning (Schaie et al., 1991). It is theorized that flexible people are more willing to play with ideas resulting in greater interest in receiving new information throughout the life-span. SLS findings suggest that older people are not becoming more rigid, rather, younger people are becoming more flexible. Many people experience noticeable ability declines in the late 60s and 70s, and it is not until the 80s that the average older adult will fall below the middle range of performance for young adults (Schaie, 2005).

Personality - Studies have found age-differences in usage of defense mechanisms and coping strategies (Rhodes & Giovannetti, 2020; Labouvie-Vief, Hakim-Larson, & Hobart, 1987; Diehl, Coyle, & Labouvie-Vief, 1996). Older adults managed their emotions through more mature defense mechanisms, while younger people reacted to psychologically demanding situations by acting-out against others, projecting their anger on others, or regressing to more primitive behavior. Older individuals were more likely to use defense mechanisms which controlled their negative emotions or that attempted to put the situation into perspective. Regarding coping, older adults were less likely to react in self-destructive or emotional ways, instead, they tried to understand the situation and find a way around it, through problem-focused coping and other strategies such as suppressing their negative feelings or channeling those feelings into productive behavior. Rhodes and Giovannetti (2017) observed strong relationships between grit (defined as having passion and perseverance for long-term goals) and multiple aspects of successful aging, including enhanced physical functioning, emotional well-being, social functioning, energy, and overall health. Similarly, Kim and Lee (2015) found that grit was more strongly associated with successful aging compared to physical functioning and socioeconomic status, implying that

elders who exhibit more grit may continue to age successfully despite lack of economic resources and restricted physical functioning given disease and disability. Additionally, older persons with higher verbal intelligence and more mature levels of ego functioning were more likely to use defense mechanisms and coping strategies utilizing thoughtful analysis, regulation of feelings, and realistic assessments and reactions to difficult situations.

Gender differences revealed that women tended to avoid unpleasant or stressful situations, blamed themselves upon things going wrong, and sought the support of others. The "feminine" coping style included showing empathy toward others and tolerating complex and ambiguous feelings in difficult situations. Men tended to externalize their feelings and use reaction formation.

Findings from trait theories of personality indicate that, as a group, older adults become less emotionally volatile, get along better with others, and accept responsibility for their actions (McCrae & Costa, 1990).

Theories of personal control, emerging from cognitive theory, suggest that personality is driven by desire to achieve control over interchanges with the environment. Two types of control processes underlie this theory (Rothbaum, Weisz, & Snyder, 1982). In *primary control*, the person's desires and goals prevail over any environmental constraints, for example, one might climb the lower shelves in a grocery store to reach the top shelf which secures a desired box of cookies. *Secondary control* involves changing the perceptions of one's goals or desires instead of the environment itself, hence, one might accept a different box of cookies within closer reach. Studies show that age-related changes in physical, cognitive, and social processes lead older adults to increasingly choose secondary control (Heckhausen, 1997). Aging individuals can avoid the frustration of inability to choose primary control by channeling their energy into attainable goals redefined through the process of secondary control.

Further, both primary and secondary control can interact with selectivity - focusing energy on a more narrow objective or changing one's approach to personal goals, or compensation - making environmental changes that will offer help in goal-attainment or downgrading the desirability of a failed goal. Older adults encounter more adverse events than younger individuals, including cognitive decline, loss of a spouse, and physical limitations which can culminate in a lower level of perceived control (Dang & Zhang, 2025). Based on compensatory control theory, people have developed a compensatory psychological and perceptual system designed to maintain a sense of order and structure when control is lessened or absent (Kay et al. 2009). This model proposes that believing the world is structured and predictable is essential because this mindset offers the basis upon which individuals can believe they can exert control over their environment and be agentic toward goals. Compensatory control theory indicates people attempt to reaffirm the foundational belief in structure/predictability in the world when their personal control is threatened, thus, they have a strong foundation upon which to reestablish a sense of

## HEALTHY AGING

control and ability to pursue their goals (Gibbs et al., 2022).

Heckhausen (1997) proposes that healthy development in later life involves finding balance among these control strategies contingent upon opportunities and constraints resulting from the aging process. Hong et al. (2021) found that sense of control is associated with better physical and psychosocial health, and improved health-promoting behaviors. The potential for primary control still prevails even as capabilities decline in later years.

Based on the MacArthur Study of Adult Development, a large national survey of roughly 3500 adults, Lachman & Weaver (1998) observed that, despite awareness of increasing life constraints, older adults (over 60) compared to people of younger ages, experienced high levels of control. They perceived their resources and potential in a positive manner instead of focusing on losses.

Erik Erikson, an optimist at heart, shared some positive qualities of older adults in the book *Vital Involvement in Old Age* (Erikson et al., 1986). The book analyzes interviews of 29 people in the Berkeley Growth Study who were studied from birth and were in their 80s during the interviews. People who rose above the infirmities and limitations of aging were identified which led to the statement, "although impairment and a certain degree of disability may be inevitable in old age, handicap and its deleterious effects on psychosocial well-being need not necessarily follow" (Erikson et al., 1986, p. 194). Everyone in the sample was not determined to overcome adversity, however, defining oneself independently of age- or illness-based limitations differentiated the two groups.

## SOCIAL FACTORS

The following social theories explain changes in social relationships occurring in late adulthood, and factors leading to successful aging.

**Role Theory** - People play different social roles throughout life, such as student, daughter, or parent, and these roles form the basis of self-concept. Each role is associated with a particular age or stage of life. Successful aging depends on the ability to accept the role changes common to the elder years.

**Age-norms** - beliefs about age-related capabilities and limitations - regulate the roles that people of different ages can play. Norms can be formally expressed (i.e., required retirement policies) or as is more common, informal. Individuals also have norms regarding the appropriateness of their own behavior at specific ages, such that "social clocks become internalized and age norms operate to keep people on the time track" (Hagestad and Neugarten, 1985).

Older adults encounter various role dilemmas: they are more likely to lose previously-held roles than to gain new ones, and many role losses are irreversible; roles become more ambiguous; and the transition from the worker to retiree role yields role discontinuity - knowledge gained at one age level may be useless or conflicting at the next age level. Role loss

can produce a decrease in social identity and self-esteem because roles are the basis of one's self-concept (Rosow, 1985). Though old age encompasses role loss, role gains can occur, such as volunteer, part-time worker, mentor, grandparent, and so on.

**Role Theory** predicts that life satisfaction declines as people grow older due to loss of roles. Roles contribute to self-image and usually each role supports self-esteem. Fewer roles equates to less support for self-esteem. Role Theory may be valid for some elders and in some societies, but many older people retain high life satisfaction and well-being, and seek role replacement, hence, this theory does not account for all factors needed for successful aging (Rosenberg, 2022).

**Activity Theory** - Assumes that active older adults will be more satisfied and better adjusted than less active elderly. The theory suggests that one's self-concept is validated through involvement with roles characteristic of middle age, hence, older people are recommended to maintain as many middle-age activities as possible, and to substitute new roles for roles lost by widowhood or retirement (Lemon, Bengston, and Peterson, 1972). To lessen society's withdrawal from the elderly, older adults must deny the existence of old age through maintaining middle-age lifestyles as long as possible, which includes remaining active, keeping busy and staying young. Behavior inappropriate to middle age is deemed to be maladaptive.

This perspective supports our society's value system which emphasizes work and productivity, and it reflects gerontological practitioners' attempts to devise new roles for older people that offer responsibilities and obligations.

Longino & Kart (1982) found that activities that socially connect people, such as having dinner with friends or engaging in hobbies through group activities had a greater tendency to improve life satisfaction than formal or solitary activities. Harlow & Cantor (1996) also determined that the social component was significant in that sharing tasks was a relevant predictor of life satisfaction, especially among retirees. Schroots (1996) noted that ability to do things despite limitations positively impacted successful aging. These studies propose that the activity type may be important instead of only the frequency of engagement.

Despite the logic of, "To be happy as you grow older, stay active," empirical support for activity theory is mixed. The theory has some inherent biases, methodological flaws, and doesn't account for many older adults with lower activity levels who remain content (Rosenberg, 2022).

**Disengagement Theory** - This controversial theory proposes that older people, undergoing losses of roles and energy, want to be freed from society's expectations of competitiveness and productivity (Bengston & Settersten, Jr., 2016; Marshall & Clarke, 2007). Disengagement is seen as adaptive behavior, facilitating self-worth and tranquility while engaging in more peripheral social roles (Adams, 2004; Birren & Schroots, 2001). In support, Cumming and Henry (1961) suggest that disengaged older people, who are released from employment

roles, participate better in family relationships than employed people. The process of disengagement for men is often abrupt as they discontinue their occupational roles, while for women it is more gradual as they transition from what is frequently their central role as parent. More employed women may experience this process differently.

Disengagement is considered functional for society as well. Cumming and Henry (1961) suggest that all societies must transfer power from older to younger generations. Retirement policies, for example, allow younger people with new energy and skills to move into occupational roles.

Critics of this theoretical perspective argue that not everyone disengages, in fact, many people in their 80s are employed, healthy, and socially active (Baltes, 1987; Neumann, 2000; Schroots, 1996). Disengagement and activity theory do not consider variability in individual preferences (Achenbaum & Bengston, 1994; Marshall, 1996; Hochschild, 1975). Not all adults want to disengage as they age; some are happy to be doing less but others are content to be doing as much or more than in the past (Rosenberg, 2022). Disengagement theory has generally lacked empirical support.

**Continuity Theory** - The weaknesses of disengagement and activity theories created a third social-psychological theory of adaptation in the elder years. Continuity theory postulates that older adults substitute similar types of roles for lost ones, and they continue to exercise typical ways of adapting to the environment to maintain psychological continuity, and continuity of social behavior and circumstances (Neugarten, Havighurst, and Tobin, 1968). Personality plays a vital role in adjusting to aging, hence, a previously active and social person is unlikely to sit quietly at home during elder years. Essentially, this theory promotes that, with age, we become more of what we were when younger. Significant personality traits and core values become even more important to the aging individual. Continuity theory believes that successful aging is a function of maintaining a mature, integrated personality while growing old, and this underscores life satisfaction (Efklides et al. 2003; Troll & Skaff, 1997; Agahi, et al. 2006; Neugarten, Havighurst, and Tobin, 1968). In this model, people rely on their own standards for successful aging instead of adjusting to a common norm.

Though continuity theory has some intuitive appeal, it also has limitations (Rosenberg, 2022). It stresses earlier stages of development as criteria for successful aging, and assumes that people attempt to maintain a certain pattern of behavior through the life span. In fact, maintaining previous patterns can be maladaptive when change is indicated (Fox, 1981-82), and releasing oneself from former roles can have positive effects (Birron & Schroots, 2001; Guttman, 1974). For example, it is considered healthy for females to adopt “masculine” personality traits with age and for men to act on tendencies that are “feminine.” It is difficult to empirically test continuity theory due to its complexity.

**The Elderly as a Subculture** - This theory believes that older adults maintain their self-concept and social identity through

membership in a subculture of older persons (Rosenberg, 2022). An aging subculture has two effects upon older persons: identification as being old, hence, being socially and culturally distant from the rest of the youth-oriented society; and a growing group consciousness that offers potential for political power and social action. Similar to viewing older adults as constituting a subculture is identifying the aged as a minority group (Blau, 1981), being discriminated against because of age.

This theory helps in understanding the role and status of older people, but the concept of a disadvantaged aging subculture does not apply to all older persons. Some elders have high status and financial security, therefore, this theory is limited in predicting behavior.

**Age Stratification Theory** - Proposes that just as societies are stratified by socioeconomic class, every society places people into categories or strata based on age, specifically, young, middle-aged, and old (Riley, 1994; Marshall, 1996; Uhlenberg, 1996; Yin & Lai, 1983; Hagestad & Dannefer, 2002; Unlenberg, 2000). Age stratification of roles frees and limits older persons, for example, older people are free from many mandatory adult roles but norms for age-appropriate behavior discourage them from working part-time or returning to school (Rosenberg, 2022).

Two factors that explain many differences in how people think, act, and contribute to society are termed the life course dimension and the historical dimension. *Life course dimension* suggests that individuals of the same chronological age or stage in the lifecycle (i.e., infancy, childhood, adolescence, early adulthood, etc.) share much in common, such as biological development, roles they have experienced, and potential years ahead, whereas people at different life stages differ in these areas. *Historical dimension* submits that people born at the same time period (cohort) share a similar historical and environmental past, present and future. Major events, such as the two World Wars, the Civil Rights Movement, space exploration, and technological advances differentially affect cohorts' values and behaviors. Often, we choose friends with similar values perhaps stemming from common experiences.

The process of *cohort flow* is illustrated by individuals stepping onto an escalator at birth (Riley, Johnson, and Foner, 1972), resulting in those who began at the bottom at the same time moving up collectively. The age group does not remain stable during the upward movement, however, as people develop unique social attributes that affect the probability of remaining on the escalator for the entire ride. Similarly, the dynamic process of age stratification confirms that different cohorts age in different ways as they “move up the escalator” (move through the age strata) due to socioeconomic factors.

Age stratification theory has been criticized for evaluating age by the too narrow concepts of chronology and life stage and not considering personal characteristics. Moreover, individual differences increase with age suggesting that cohorts become less cohesive with age rather than more unified. Age stratification can illuminate how society uses

age to fit people into structural niches in the social world.

**Interactionist Perspectives** - Emphasize the importance of dynamic interaction between older people and their social world (Rosenberg, 2022). Given change, older individuals are advised to try to master the changing situation while utilizing resources within their environment to preserve a positive self-concept.

The *symbolic interactionist* view of aging indicates that the interaction of the environment, individuals, and their encounters in the environment affect the type of aging process people experience (Gubrium, 1973). For example, an older person who becomes confused after moving to a new dwelling may be labeled as senile, when simply lowering the stress of moving could reduce the confusion. Creating new alternatives is recommended, for example, withdrawal from social involvement is not inevitable with aging, rather, it is one possible outcome of a person's interaction which can be changed.

*Labeling theory* states that people gain self-concept from interacting with others in their social milieu. The labeling process begins, for example, when someone ready to retire acts non-productively because others have defined retirement in this manner.

The *social breakdown theory* argues that older people who accept negative labeling begin to act in dependent ways and their independence diminishes.

The *social reconstruction model* believes that even small changes in restructuring the environment can improve the quality of life for elders. Unfortunately, an assumption within this model is that socioeconomic class creates inequality which limits one's potential mastery over life (Tindale and Marshall, 1980).

**Social Exchange Theory** - States that a vital factor in defining the elderly's status is the balance between their societal contributions, which are determined by their control of power resources, and the costs involved in supporting them (Rosenberg, 2022). Given possession of material goods, abilities, achievements, and other qualities deemed desirable by the culture, individuals can exert power in their social relationships. Those elderly in possession of fewer power resources than younger people may experience diminished status.

This theory proposes that older people disengage because they have little of value to exchange, in turn, they are often forced to accept the retirement role. Inability to work limits their access to two valuable power resources - material possessions and authority positions.

Dowd (1975) asserts that the only major power source unscathed by aging is the category of generalized reinforcers, such as respect, approval, recognition and support; this category is less valued than other resources, which limits elderly in influencing exchange rates. Despite limited resources, most older adults attempt to exercise a degree of reciprocity and active control in the management of their lives.

Additional empirical research is needed to assess the value of exchange theory as explanation for the aging process.

**Political Economy of Aging** - This perspective believes that social class is a barrier to older people's access to important resources, and that powerful societal groups maintain their own interests by perpetuating class inequalities (Rosenberg, 2022; Marshall & Clarke, 2007; Minkler and Estes, 1984; Walker, 1981). Estes asserts that social, political, and economic conditions affect how the elderly's concerns are defined and treated. The elderly's problems are thought to be socially constructed due to societal conceptions of aging and the aged. The aging process itself is not the problem, instead, the issue is the societal conditions encountered by older people lacking sufficient income, health care, or housing, which are needs created by a capitalist society. Estes argues that major policy changes are needed which would not separate the elderly because of their age, and would improve social perceptions and objective conditions of the aged.

Essentially, the political economy of aging model analyzes sociopolitical factors that separate social policies for groups, such as the elderly. The main limitation of this critique is lack of empirical research.

## SOCIAL SUPPORT SYSTEMS

The *socioemotional selectivity theory* indicates that throughout adulthood, people decrease the range of their relationships in order to maximize social and emotional gains and minimize risks (English & Carstensen, 2014; Perkins, et al., 2012; Carstensen, 1987). Further, as individuals grow older, they are more focused upon maximizing emotional rewards and less interested in seeking information or knowledge through their relationships. This shift occurs as people become increasingly aware that they are "running out of time." Increasing motivation for emotionally meaningful goals can be beneficial, for example, when responding to irrecoverable losses and missed opportunities (Strough et al., 2008; 2019), but caution is advised to avoid such emotional well-being focus from preventing careful deliberation of the facts (Mikels et al., 2013). Carstensen (2006) noted the recognition of life's finitude motivates pursuit of present-oriented, emotionally meaningful positive experiences as opposed to future-oriented information seeking. This mindset was found to increase choices by research participants for options that promoted calmness and reduced regret (Jiang et al., 2016; Strough et al., 2019). Young adults feeling real or artificial time constraints show similar desire toward emotional gains from their social interactions as older adults (Carstensen, Isaacowitz, & Charles, 1999). In other words, at this stage, we want to spend time with those who have been closest to us rather than establishing new friends or acquaintances.

Older adults are often better able to control expression of their emotions (Lawton, Kleban, Rajagopal, & Dean, 1992), and they have fewer negative emotions, possibly because of greater ability to regulate their affective experiences (Gross et

al., 1997). Older adults display heightened memory and attention for positive information in contrast to negative information, which is termed the positivity effect (Carstensen & DeLiema, 2018). For example, older adults viewed and remembered a greater amount of positive rather than negative information compared to younger adults when making hypothetical health-care decisions (Lockenhoff & Carstensen, 2007; Depping & Freund, 2013). Relative to younger adults, older adults exhibited less rumination about losses which contributed to their reporting more positive and less negative emotions after an experimental randomly determined loss on a gambling task (Bruine et al., 2018). These age differences in emotions support the importance of emotional experience in older adults and imply that they would prefer to spend time with their long-term relationship partner.

Over 50% of those over age 65 are married and living with a spouse in an independent household (U.S. Senate Special Committee on Aging, 1992). Due to women's longer life expectancy and fewer options for re-marriage, 40% of women age 65 and over are married, as compared to 74% of men. Women represent 80% of the elderly who live alone.

Marital tensions due to changing roles and expectations may be heightened for older couples as partners change roles through retirement, post-parenthood, or illness. Inability to negotiate role expectations and consequent disagreements can lead to feelings of inequity and depression in older spouses (Holahan, 1984). Despite such potential tension, most older partners seem satisfied, with men being more satisfied with marriage and their emotional need fulfillment than women (Chappell, 1990; Gilford, 1984).

Research indicates that there is no physiological reason for a lessening of older women's ability to enjoy sexuality. In fact, the main obstacle to a woman's ability to experience a rewarding sex life after age 60 is the unavailability of a partner (Marsiglio & Donnelly, 1991; Masters & Johnson, 1966; Pfeiffer, Verwoerd, & Davis, 1974). Regarding men, the ability to have an erection may be affected by various physiological and psychological factors from middle age onward. Men over age 60 experience normal age-related changes in sexual functioning involving a slowing down and decrease of intensity in their movement through the sexual response cycle. Arousal is slower, orgasm is shorter, and less seminal fluid is ejected (Masters & Johnson, 1970). Roughly 15-25% of men 65 and over have erectile dysfunction (National Institute of Diabetes and Digestive and Kidney Diseases, 1999; National Institutes of Health, 1992). Health can significantly affect erectile functioning, especially disorders of the cardiovascular and endocrine systems. Stress, depression and anxiety can also interfere with sexual functioning.

The majority of older adults have positive and accepting attitudes toward sexuality and believe that it has importance in their lives (Starr & Weiner, 1981). Findings from this survey of 800 participants over age 60 obtained through community centers indicated: 76% expressed that sex has a positive effect on their health; 75% felt that sex feels the same or better compared to their younger days, with 41% of women and

27% of men stating that sex is better now; 99% would like to have sex if it were available; and 91% approved of sex and living together without marriage for older adults.

Estimation of sexual activity in single, divorced, and widowed people ranges from 32% (Starr, 1985) to over 50% of women and 75% of men in their 60s and 70s (Brecher, 1984).

Older people who live alone without social support systems have higher probability of institutionalization, lower personally reported well-being, and greater difficulty adjusting to widowhood compared to those with strong social supports (Lopata, 1979; Kasper, 1988; Wallston et al., 1983).

About 80% of people over age 75 with functional disabilities and living in the community are assisted by their families (U.S. Senate Special Committee on Aging, 1988). Wives and daughters perform most of the personal care (Cantor, 1991; Stoller, 1990), in fact, women represent over 80% of the family caregivers to chronically ill elderly. Without such family support, many of these elders would be institutionalized and the number of nursing home residents would triple (Brody, 1985). Those without family ties, mainly widowed women and the very old who have outlived family members, have the greatest likelihood of institutionalization.

With age, siblings frequently renew past ties, forgive conflict and rivalry, and become closer, often through shared reminiscence (Cicirelli, 1991; Brubaker, 1990).

After one's spouse, adult children represent the most relevant source of support and social contact for older adults. The majority of those over age 65 live near their children but do not share the same home; most older persons report not wanting to live with their children for reasons of privacy and autonomy. Less than 20% of elderly live in their children's households, but this number increases to 33% of all men, and 50% of all women age 65 and over who are widowed, separated, or divorced that live with their children or other family members. Roughly 80% of older adults with children live less than one hour away from at least one child, and over 75% have phone discussion at least weekly (AARP, 1991; Crimmins and Ingegneri, 1990; Shanas, 1979, 1980).

Approximately 30% of people over 65 and 45% of those 85 and over live alone, and they are most likely to be women, ethnic minorities, the oldest-old, and of low socioeconomic status. These individuals often rely more on community services than on friends and neighbors for support to continue independent living; and they report lower life-satisfaction than married people (Kasper, 1988).

Several intervention models are designed to increase peer group interaction for the elderly's well-being. *Personal network building* consists of caring and concerned neighbors, for example, who act as "natural helpers" by offering support or services to elderly in need. Further, "gatekeepers," such as newspaper or postal carriers (by observing whether the newspaper or mail is brought in) can connect isolated elderly with sources of assistance as Senior Information and Assistance lines. Churches may also serve as personal networks by providing a surrogate family for elders or facilitating church members to offer various types of

assistance.

*Volunteer linking* utilizes volunteers to perform chores, offer peer counseling and serve as Friendly Visitors.

*Mutual help networks* offer problem-solving and reciprocal exchange of resources, for example, groups focusing upon shared problems such as widow-to-widow programs or stroke clubs can offer new skills, expanded social networks, and greater solution capabilities.

Another intervention model is *neighborhood and community development* which strives to maximize a community's self-help and problem-solving capacities. Older adults in a low-income neighborhood, for example, can unite as a "family" to solve personal and community concerns.

Social relationships maintain importance throughout life. A biopsychosocial model of adult relationships is termed the *social convoy* which suggests that people possess a network of close relationships that "carries" them or offers social support throughout their lives (Kahn & Antonucci, 1980). Though the convoy may change, it maintains its size of 5 to 10 close social ties and the exchange of support within the social network. The model resembles a set of concentric circles in which the closest family members reside within the innermost circles and more distant relationships extend outward (Antonucci & Akiyama, 1987; Chatters, Taylor, & Jackson, 1986). Support is exchanged in both directions from and to others within the convoy, and despite older adults tending to feel they are receiving more than providing support, exchange relationships remain reciprocal through adulthood (Antonucci & Akiyama, 1987).

The degree of feeling satisfied with the number of people in the closest circle is a predictive factor in one's well-being (Antonucci et al., 2010; Antonucci, Fuher, & Dartigues, 1997). The structure, quality, and function of a person's social convoy dynamically and continually evolves given the individual's personal and situational characteristics across their lifespan (Fuller, et al., 2020; Koehly & Manalel, 2023). Satisfaction with the social network may contribute to improved physical functioning and health in the cardiovascular, endocrine, and immune systems (Yang et al., 2016; Uchino et al., 2018; Uchino, Cacioppo, & Kiecolt-Glaser, 1996).

## THE NATURE OF HEALTHY AGING

Aging researchers study factors which contribute to the survival capability of older adults, whereas theorists of healthy aging examine such concepts as "mental health" factors which allow older adults to transcend losses and physical limitations often accompanying later years.

Findings from a research project called the McArthur Foundation Study of Aging in America found the following three factors extant in successful aging (Rowe & Kahn, 1998; Rowe, 2018): a) absence of disease and the disability associated with disease, and not displaying risk factors that increase likelihood of disease and disability; b) revealing high cognitive and physical functioning which enhances potential of being active and competent; and c) showing "engagement

with life" which means involvement with productive activity and interacting with people.

Supportive of the possibility for healthy aging are the findings that most older people do not become depressed, personality development in later life moves toward enhanced adaptiveness, and Erikson's theory of psychosocial development suggests that elders' level of development is fuller than that of younger people whose egos are yet to be thoroughly tested by time and experience.

The *paradox of well-being* suggests that the majority of older people experience relatively high levels of well-being (Mroczek & Kolarz, 1998; Swift et al. 2014; Hansen & Slagsvold, 2012). Older adults maintain a positive view of themselves and their life situations despite hardships and this favorable perspective seems to be the norm. A study of over 32,000 Americans from 1972 to 1994 revealed that the large majority of older individuals rated themselves as "very" or "pretty" happy (Mroczek & Kolarz, 1998). Research from many countries supports this concept of increased feelings of satisfaction with aging (Diener & Suh, 1998; Hansen & Slagsvold, 2012).

Older adults often utilize various methods to maintain a positive perspective, such as adaptation, coping mechanisms, including problem-focused and emotion-focused strategies, *social comparison* - assessing negative circumstances of others and realizing one's own situation could be worse (Michalos, 1985; Weiss et al., 2025), and the process of interpreting life events in a positive manner.

Beyond maintaining a positive mental attitude, older adults can experience a renewal in productivity and creativity culminating in a secondary peak, and those with high creative potential are more likely to exhibit a high rate of productivity both early and late in their careers (Simonton, 1998; Kozbelt, 2008). Lehman (1953), and Kozbelt (2008) noted that "older thinkers" have produced great achievements as evidenced by an upturn in artists producing "best" paintings in their 70s. For example, Michaelangelo contributed the "Pieta Rondanini" at age 89, similarly, Benjamin Franklin invented bifocals at 78, and Galileo contributed greatly to the field of mechanics at 74. Such contributions may be motivated by closeness to death, urge to leave a legacy, or as reaction to age-related changes or health problems. Retirement, with the right frame of mind, can open new doors to enjoyment and productivity.

Abud et al. (2022) searched five electronic databases (EMBASE, MEDLINE, Cochrane, PsychInfo, and CINAHL) from 2010 to 2020 to identify contemporary evidence on factors that can potentially be determinants of healthy aging. The ten identified ten determinants for healthy aging included physical activity, diet, self-awareness, outlook/attitude, life-long learning, faith, social support, financial security, community engagement, and independence. Results showed an increasing acknowledgement of the instrumental role of social and mental/cognitive well-being as significant determinants of healthy aging.

Centenarians are those who live between ages 100 and 109, and in the United States, they numbered roughly 82,000 in

year-2016, 92,000 in year-2020, with a projected 500,000 in year-2060 (Duffin, 2020). The increase is attributed to improvement in vaccines, antibiotics, hygiene, and sanitation; genetics has also been a factor (Fessenden, 2020). The U.S. Census report of 2010 indicates centenarians were predominantly Caucasian women living in urban areas of the Southern states. Approximately 43% of the male centenarians lived with family members while females most often lived in nursing homes. The countries with the largest number of centenarians (in decreasing order) are Japan, America, and France (Aging Analytics Agency, with Gerontology Research Group, 2020).

Research shows many long-lived people do not smoke, are not obese, manage stress well, and are less likely to develop various chronic diseases such as hypertension, heart disease, cancer, or diabetes (NIH, 2020; Sebastiani et al., 2013).

Super-centenarians are those who live until at least 110 years, and there are 300-450 alive at any time worldwide. Analysis of 32 super-centenarians in the United States observed that "A surprisingly substantial portion of these individuals were still functionally independent or required minimal assistance" (Schoenhofen et al., 2006). Most functioned independently until after age 100, with no indications of frailty until approximately age 105. It is concluded these unusual individuals are long-lived for "rare and unpredictable" reasons (Willcox et al., 2008). The number of super-centenarians is predicted to grow as future centenarians live longer due to healthier lifestyle (Robine & Vaupel, 2001).

A growing emphasis on holistic health has broadened definitions of health and wellness. Currently, wellness involves the individual's whole being, including physical, emotional, mental, and spiritual. Wellness includes attaining a balance between one's internal and external environment along with one's emotional, spiritual, social, cultural, and physical processes. The individual strives to achieve wellness through effective health practices (Touhy & Jett, 2022).

The process of achieving wellness requires work, and plateaus may be reached during the ascension to higher-level wellness. Even chronic illnesses, acute events, and crises can be viewed as potential stimulus for growth and a return to traversing the wellness continuum.

Health in later life often coincides with functional ability (i.e., ability to do what is deemed important by the person) instead of the absence of disease (Touhy & Jett, 2022).

A holistic perspective of health emphasizes strengths, resilience, resources, and capabilities as opposed to searching for evidence of potential pathological conditions. A wellness viewpoint believes that everyone has an optimal level of health independent of their situation or functional ability (Touhy & Jett, 2022).

## REFERENCES

- Abud, T., Kounidas, G., Martin, K.R., et al. (2022). Determinants of healthy ageing: A systematic review of contemporary literature. *Aging Clinical Experimental Research*, 34, 1215-1223.
- Adams, K.B. (2004). Changing investment in activities and interests in elders' lives: Theory and measurement. *International Journal of Aging & Human Development*, 58(2), 87-108.
- Adelman, R.C. (1980). Hormone interaction during aging. In R.T. Schimke (Ed.), *Biological mechanisms in aging*, p.686. Washington, DC: U.S. Department of Health and Human Services.
- Allen, M.F.S. (1997). *Aging and human longevity*. Boston, MA: Birkhauser.
- Agahi, N., Ahacic, K., & Parker, M.G. (2006). Continuity of leisure participation from middle age to old age. *The Journals of Gerontology*, 61B(6), S340-S346.
- Aging Analytics Agency (with the Gerontology Research Group; 2020): *Supercentenarians Landscape Overview. Top-100 Living, top-100 longest lived, top-25 socially and professional active*.
- Ahmad, F.B., Cisewski, J.A., Anderson, R.N. (2024). Leading causes of death in the US, 2019-2023. *JAMA*, 332(12): 957-958.
- Ahmed, B., Rahman, A.A., Lee, S., & Malhatra, R. (2024). The implications of aging on vascular health. *International Journal of Molecular Science*, 25(20), 11188.
- Alamro, H., Thafar, M.A., Albaradei, S. et al. (March 27, 2023). Exploiting machine learning models to identify novel Alzheimer's disease biomarkers and potential targets. *Scientific Reports*, 13, Article number: 4979.
- Aldwin, C.M., Gilmer, D.F. (2004). *Health, illness and optimal aging: Biological and psychological perspectives*. Thousand Oaks, CA: Sage Publications.
- Almutairi, K., Nossent, J., Preen, D. et al. (2021). The global prevalence of rheumatoid arthritis: a meta-analysis based on a systematic review. *Rheumatology International*, 41, 863-877
- American Academy of Dermatology Association (2025). Sunscreen faqs. [www.aad.org/media/stats-sunscreen](http://www.aad.org/media/stats-sunscreen)
- American Federation for Aging Research. (2016). *Biology of Aging*. [www.afar.org/infoaging](http://www.afar.org/infoaging)
- American Association of Retired Persons (AARP). *A profile of older Americans*, 1990. Washington, D.C.: AARP, 1991.
- Antonucci, T.C., & Akiyama, H. (1987). Social networks in adult life and a preliminary examination of the convoy model. *Journal of Gerontology*, 42, 519-527.
- Antonucci, T.C., Fiori, K.L., Birditt, K., & Jackey, L.M.H. (2010). Convoys of social relations: Integrating life-span and life-course perspectives. In M.E. Lamb & A.M. Freund (Eds.), *The Handbook of Life-Span Development* (Vol. 2, pp. 434-473. John Wiley & Sons, Inc.
- Antonucci, T.C., Fuhrer, R., & Dartigues, J. (1997). Social relations and depressive symptomatology in a sample of community-dwelling French older adults. *Psychology and Aging*, 12, 189-195.
- Arking, R. (1991). *Biology of aging: Observations and Principles*. Englewood Cliffs, NJ: Prentice-Hall, Inc.
- Asejeje, F.O., & Ogunro, O.B. (March, 2024). Deciphering the mechanisms, biochemistry, physiology, and social habits in the process of aging. *Archives of Gerontology and Geriatrics Plus*, Volume 1, Issue I.
- Bacon, C.G, Mittleman, M.A., Kawachi, I. et al. (2003). Sexual function in men older than 50 years of age: results from the health professionals follow-up study. *Annals of Internal Medicine*, Vol. 139:pp. 161-8.
- Baddely, A.D. (1986). *Working memory*. London: Oxford University Press.
- Baltes, P.B. (1987). Theoretical propositions of life-span developmental psychology: On the dynamics between growth and decline. *Developmental Psychology*, 23, 611-626.
- Baltes, P.B., & Graf, P. (1996). Psychological aspects of aging: Facts and frontiers. In D. Magnusson (Ed.), *The lifespan development of individuals: Behavioral, neurobiological, and*

- psychosocial perspectives* (pp. 427-460). New York: Cambridge University Press.
- Bays, H.E. et al. (2022). Ten things to know about ten cardiovascular disease risk factors - 2022. *American Journal of Preventive Cardiology*, Vol 10, June, 2022.
- Besedes, T., Deck, C., Sarangi, S., & Shor, M. (2012). Age effects and heuristics in decision making. *Review of Economics and Statistics*, 94, 580-595.
- Bengston, V., & Schaie, K.W. (1999). *Handbook of theories of aging*. New York: Springer Publishing Company.
- Bengston, V.L., & Settersten Jr., R.A. (2016). *Handbook of Theories of Aging*, 3rd ed. Springer Publishing Company.
- Besedes, T., Deck, C., Sarangi, S., & Shor, M. (2012). Age effects and heuristics in decision making. *Review of Economics and Statistics*, 94, 580-595.
- Birren, J.E., & Schroots, J.J.F. (2001). History of gero-psychology. In J.K.E. Birren (Ed.), *Handbook of psychology of aging* (5th ed., pp. 3-28. San Diego, CA: Academic Press.
- Bjorntorp, P. (1996). The regulation of adipose tissue distribution in humans. *International Journal of Obesity and Related Metabolic Disorders*, 20, 291-302.
- Blanchflower, D.G. (2020). Is happiness U-shaped everywhere? Age and subjective well-being in 145 countries. *Journal of Population Economics*, Vol. 34, pp. 575-624.
- Blau, Z. (1981). *Aging in a changing society* (2nd ed.). New York: Franklin Watts.
- Boonen, S., Lesaffre, E., Dequeker, J., Aerssens, J., Nijs, J., Pelemans, W., & Bouillon, R. (1996). Relationship between baseline insulin-like growth factor-I (IGF-I) and femoral bone density in women aged 70 years. Potential implications for the prevention of age-related bone loss. *Journal of the American Geriatrics Society*, 44, 1301-1306.
- Botwinick, J. (1977). Intellectual abilities. In J.E. Birren & K.W. Schaie (Eds.), *Handbook of the psychology of aging* (pp. 580-605). New York: Van Nostrand Reinhold.
- Bouchard, J., & Villeda, S.A. (2015). Aging and brain rejuvenation as systemic events. *Journal of Neurochemistry*, 132.1, 5-19.
- Brecher, E. (1984). *Love, sex, and aging*. Boston: Little, Brown.
- Breit, M., Scherrer, V., Tucker-Drob, E. M., & Preckel, F. (2024). The stability of cognitive abilities: A meta-analytic review of longitudinal studies. *Psychological Bulletin*, 150(4), 399-439.
- Brody, E. (1985). Parent care as a normative family stress. *The Gerontologist*, 25, 19-30.
- Brookmeyer, R., & Kawas, C. (1998). Projections of Alzheimer's disease in the United States and the public health impact of delaying disease onset. *American Journal of Public Health*, 88, 1337-1342.
- Brubaker, T.H. (1990). Families in later life: A burgeoning research area. *Journal of marriage and the family*, 52, 959-981.
- Bruine de Bruin, W., Van Putten, M., Van Emden, R., & Strough, J. (2018). Age differences in emotional responses to monetary losses and gains. *Psychology and Aging*, 33, 413-418.
- Bruine de Bruin, W., Strough, J., & Parker, A. M. (2014). Getting older isn't all that bad: Better decisions and coping when facing "sunk costs." *Psychology and Aging*, 29, 642-649.
- Bruine de Bruin, W., Van Putten, M., Van Emden, R., & Strough, J. (2018). Age differences in emotional responses to monetary losses and gains. *Psychology and Aging*, 33, 413-418.
- Bugg, J.M., DeLosh, E.L., Davalos, D.B., & Davis, H.P. Age differences in Stroop Interference: Contributions of general slowing and task-specific deficits. *Aging Neuropsychological Cognition*, 14: 155-167.
- Burke, D.M. (1997). Language, aging and inhibitory deficits: Evaluation of a theory. *Journal of Gerontology: Psychological Sciences*, 52B, p. 254-264.
- Burke, D.M., & Shafto, M.A. (2004). Aging and language production. *Current Directions in Psychological Science*, 13, 21-24.
- Burraco, A.B., Fernandez, Lopez, I., Fernandez-Perez, M. & Ivanova, O. (2025). *Biolinguistics at the cutting edge: Promises, achievements, and challenges*. Walter de Gruyter, Berlin/Boston.
- Cantor, M. (1991). Family and community: Changing roles in an aging society. *The Gerontologist*, 31, 337-340.
- Carstensen, L.L. (1987). Age-related changes in social activity. In L.L. Carstensen & B.A. Edelman (Eds.), *Handbook of clinical gerontology* (pp. 222-237). Elmsford, NY: Pergamon Press.
- Carstensen, L. L. (2006). The influence of a sense of time on human development. *Science*, 312, 1913-1915.
- Carstensen, L.L (2021). Socioemotional Selectivity Theory: The Role of Perceived Endings in Human Motivation. *The Gerontologist*, Volume 61, Issue 8, pp. 1188-1196
- Carstensen, L. L., & DeLiema, M. (2018). The positivity effect: A negativity bias in youth fades with age. *Current Opinion in Behavioral Sciences*, 19, 7-12.
- Carstensen, L.L., Isaacowitz, D.M., & Charles, S.T. (1999). Taking time seriously: A theory of socioemotional selectivity. *American Psychologist*, 54, 165-181.
- Cattell, R.B. (1963). Theory of fluid and crystallized intelligence: A critical experiment. *Journal of Educational Psychology*, 54, 1-22.
- Cattell, R.B. (1971). *Abilities: Their structure, growth, and action*. Boston: Houghton Mifflin.
- Cavanaugh, J.C., & Green, E.E. (1990). I believe, therefore I can: Self-efficacy beliefs in memory aging. In E.A. Lovelace (Ed.), *Aging and cognition: Mental processes, self-awareness, and interventions* (pp. 189-230). Amsterdam: North Holland.
- CDC/National Center for Health Statistics (2024). Mortality in the United States, 2022.
- CDC/National Center for Health Statistics (2025). Mortality in the United States, 2023.
- Cerella, J., Poon, L.W., & Williams, D.M. (1980). Age and the complexity hypothesis. In L.W. Poon (Ed.), *Aging in the 1980s* (pp. 332-340). Washington, DC: American Psychological Association.
- Chappell, N.C. (1990). Aging and social care. In R.B. Binstock and L.K. George (Eds.), *The handbook of aging and the social sciences* (3d ed.). New York: Academic Press.
- Chatters, L.M., Taylor, R.J., & Jackson, J.S. (1985). Size and composition of the informal helper networks of elderly blacks. *Journal of Gerontology*, 40, 605-614.
- Chen, H., Chen, C., Spanos, M. et al.(2022). Exercise training maintains cardiovascular health: signaling pathways involved and potential therapeutics. *Signal Transduction and Target Therapy Vol. 7*, p. 306.
- Chen, X., Hertzog, C. & Park, D.C. (2017). Cognitive predictors of everyday problem solving across the lifespan. *Gerontology*, 63(4).
- Choi, J, Lee, S.H., & Yu, S. (2024). Exploring factors influencing caregiver burden: A systematic review of family caregivers of older adults with chronic illness in local communities. *Healthcare*, 12(10), p. 1002.
- Christiansen, J.L., & Grzybowski, J.M. (1993). *Biology of aging*. Saint Louis, MO: Mosby Year Book, Inc.
- Cicirelli, V.G. (1991). Siblings as caregivers in middle and old age. In J. Dwyer and R. Coward, (Eds.), *Gender, families and elder care*. Newbury Park, CA: Sage.
- Clark, W.R. (2002). A means to end: *The biological basis of aging and death*. Oxford University Press.
- Cohen-Mansfield, J., Hazan, H., Lerman, Y.K. & Shalom, V. (2016). Correlates and predictors of loneliness in older adults: A review of quantitative results informed by qualitative insights *International Psychogeriatrics*, Vol. 28(4), pp. 557-576.
- Craik, F.I.M. (1994). Memory changes in normal ageing. *Current Directions in Psychological Science*, 3, 155-158.
- Crimmins, E.M., & Ingegneri, D.G. (1990). Interaction and living arrangements of older parents and their children: Past trends, present determinants, future implications. *Research on Aging*, 2, 3-35.
- Cumming, E., & Henry, W.E. (1961). *Growing old*. New York:

- Basic Books.
- Dang, Q., & Zhang, B. (2025). Sense of control and aging stereotyping: the role of need for structure. *BMC Psychology*, Vol. 13, Article No. 732.
- De la Fuente, M. (2013). The immune system, a marker and modulator of the rate of aging. *Immunology of aging*. Springer Berlin Heidelberg. 3-23.
- Del Missier, F., Mäntylä, T., & Bruine, deBruin, W. (2012). Decision-making competence, executive functioning, and general cognitive abilities. *Journal of Behavioral Decision Making*, 25, 331–351.
- Del Missier, F., Mäntylä, T., Hansson, P., et al. (2013). The multifold relationship between memory and decision making: An individual-differences study. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 39, 1344–1364.
- Depping, M. K., & Freund, A. M. (2013). When choice matters: Task-dependent memory effects in older adulthood. *Psychology and Aging*, 28, 923–936.
- Diamond, M.C. (1993). An optimistic view of the aging brain. *Generations*, 17, 31-33.
- Diaz-Canestro, C., Pentz, B., Sehgal, A., & Montero, D. (2022). Sex differences in cardiorespiratory fitness are explained by blood volume and oxygen carrying capacity. *Cardiovascular Research*, Vol. 118, Issue 1, pp. 334-343.
- Diaz, M.T., Avery, A.R. & Zhuang, J. (2016). The neural language systems that support healthy aging: Integrating function, structure, and behavior. *Language and Linguistics Compass* 10(7). 314-334.
- Diehl, M., Coyle, N., & Labouvie-Vief, G. (1996). Age and sex differences in coping and defense across the life span. *Psychology and Aging*, 11, 127-139.
- Diener, E., & Suh, E. (1998). Age and subjective well-being: An international analysis. *Annual Review of Gerontology and Geriatrics*, 17, 304-324.
- Digiovanna, A.G. (1994). *Human aging: Biological perspectives*. New York: McGraw Hill.
- DiLoreto, R., & Murphy, C.T. (Oct. 13, 2017). The cell biology of aging. *Molecular Biology of the Cell*, Vol. 26, No. 25.
- Dixon, R.A., & Hultsch, D.F. (1999). Intelligence and cognitive potential in late life. In J.C. Cavanaugh & S.K. Whitbourne (Eds). *Gerontology: Interdisciplinary perspectives* (pp. 213-237). New York: Oxford University Press.
- Donato, A.J., Machin, D.R., & Lesniewski, L.A. (2018). Mechanisms of dysfunction in the aging vasculature and role in age-related. *Dis Circ Res*, 123:825–848.
- Dowd, J.J. (1975). Aging as exchange: A preface to theory. *Journal of Gerontology*, 30, 584-594.
- Duffin, E. (March 6, 2020). *Number of centenarians in the U.S. 2016-2060*. <https://www.statista.com/statistics/996619/number-centenarians-us/>
- Eberhardt, W., Bruine de Bruin, W., & Strough, J. (2018). Age differences in financial decision making: The benefits of more experience and less negative emotions. *Journal of Behavioral Decision Making*, 32, 79–93.
- Edwards, M., & Abadie, L. (April 11, 2019). *NASA's Twins Study Results Published*. <https://www.nasa.gov/humans-in-space/nasas-twins-study-results-published-in-science-journal>
- Efklides, A., Kalaizidou, M., & Chankin, G. (2003). Subjective quality of life in old age in Greece; The effect of demographic factors, emotional state, and adaptation to aging. *European Psychologist*, 8, 178-191.
- English T., & Carstensen, J.L. (2014). Selective Narrowing of Social Networks Across Adulthood is Associated With Improved Emotional Experience in Daily Life. *International Journal of Behavior Development*. March 1;38(2):195-202.
- Enserink, M. (1998). First Alzheimer's disease confirmed. *Science*, 279, 2037.
- Erceg, N., Micic, M., Forouzan, E., & Knezevic, N.N. (2025). The role of cortisol and dehydroepiandrosterone in obesity, pain, and aging. *Diseases*, 13(2): 42.
- Erikson, E.H., Erikson, J.M., & Kivnick, H.Q. (1986). *Vital involvement in old age*. New York: W.W. Norton.
- Everitt, T., Davies, M., Wadsworth, L., & Yantha, B. (2023). *Nutrition in Aging*, Chapter 2, In *Theories of Aging*.
- Fang, Y., Gong, A.Y., Haller, S.T., Dworkin, L.D., Liu, Z., & Gong, R. (2020). The ageing kidney: Molecular mechanisms and clinical implications. *Ageing Research Review*, 63, 101151.
- Fernandez-Ballesteros, R., Benetos, A., & Robine, J.M. (2019). *Cambridge Handbook of Successful Aging*. Cambridge University Press.
- Fishburn, M.J., & Lateur, B.J. (1996). Rehabilitation. In D.B. Reuben, T.T. Yoshikawa, & R.W. Besdine (Eds.), *Geriatrics review syllabus* (3rd ed., pp. 93-103). Dubuque, IA: Kendall/Hunt.
- Fitch, A.K., & Bays, H.E. (2022). Obesity definition, diagnosis, bias, standard operating procedures (SOPs), and telehealth: An Obesity Medicine Association (OMA) Clinical Practice Statement (CPS). *Obesity Pillars*, Vol. 1, Article 100004.
- Formosa, M. (2023). Five decades of older adult learning: Achievements and challenges. *International Journal of Education and Ageing*, Vol. 5, No. 3, 91-104.
- Fox, J.H. (1981-82). Perspectives on the continuity perspective. *International Journal of Aging and Human Development*, 14, 97-115.
- Frette, C., Barrett-Connor, E., & Clausen, J.L. (1996). Effect of active and passive smoking on ventilatory function in elderly men and women. *American Journal of Epidemiology*, 143, 757-765.
- Fuller, H.R., Ajrouch, K.J., & Antonucci, T.C. (2020). Original voices the convoy model and later-life family relationships. *Journal of Family Theory and Review*, p. 6050.
- Gatto, L., Hallen, G., Brown, A., & Samman, S. (1996). Ascorbic acid induces a favorable lipoprotein profile in women. *Journal of the American College of Nutrition*, 15, 154-158.
- Giaquinto, A.N. (October 1, 2024). Breast cancer statistics 2024. *CA: A Cancer Journal for Clinicians*.
- Gibbs, W.C., Kim, H.S., Kay, A.C., Sherman, D.K. (Dec. 16, 2022). Who needs control? A cultural perspective on the process of compensatory control. *Social and Personality Psychology Compass*.
- Gilford, R. (1984). Contrasts in marital satisfaction throughout old age: An exchange theory analysis. *Journal of Gerontology*, 39, 325-333.
- Gluck, J., Jager, L., Auer-Spath, I., & Harbig, I.A. (2025). Finding the balance to quiet the striving: The difference between successful aging and wise aging. *The Gerontologist*, Vol. 65, Issue 1, gnae126.
- Gonzales, M.M., Kraig, E., Orr, M.E. (2022). Biological aging processes underlying cognitive decline and neurodegenerative disease. *The Journal of Clinical Investigation*. 132, (10).
- Gorbulnova, V., Seluanov, A., & Pereira-Smith, O.M. (2002). Expression of human telomerase (hTERT) does not prevent stress-induced senescence in normal human fibroblasts but protects the cells from stress-induced apoptosis and necrosis. *Journal of Biological Chemistry*, Vol. 277, Issue 41, pp. 38540-38549.
- Greene, V.L., & Ondrich, J.I. (1990). Risk factors for nursing home admissions and exits. *Journals of Gerontology*, 45, S250-S258.
- Griffin, W.S., Sheng, J.G., Royston, M.C., Gentleman, S.M., McKenzie, J.E., Graham, D.I., Roberts, G.W., & Mrak, R.E. (1998). Glial-neuronal interactions in Alzheimer's disease: the

- potential role of a “cytokine cycle” in disease progression. *Brain Pathology*, 8, 65-72.
- Gross, J.J., Carstensen, L.L., Pasupathi, M., Tsai, J., Goettestam Skorpen, C., & Hsu, A.Y.C. (1997). Emotion and aging: Experience, expression, and control. *Psychology and Aging*, 12, 590-599.
- Grote, C., Reinhardt, D., Zhang, M., & Wang J. (2019). Regulatory mechanisms and clinical manifestations of musculoskeletal aging. *Journal of Orthopaedic Research*, 37:1475–1488.
- Grundey, S.M. et al. (2019). Guideline on the management of blood cholesterol: A report of the American College of Cardiology/Heart Association task force on clinical practice guidelines. *Journal of the American College of Cardiology*. Vol. 73, Issue 24, pp e285-350.
- Gubrium, J.F. (1973). *The myth of the golden years*. Springfield, IL: Charles C. Thomas.
- Gunes, S., Aizawa, Y., Sugashi, T., et al. (April 29, 2022). Biomarkers for Alzheimer's Disease in the current state: A narrative review. *International Journal of Molecular Science*, 23(9): 4962.
- Gunes, S., Aizawa, Y., Sugashi, T., et al. (April 29, 2022). Biomarkers for Alzheimer's Disease in the current state: A narrative review. *International Journal of Molecular Science*, 23(9): 4962.
- Guo, J., Huang, X., Dou, L., Yan, M., Shen, T., Tang, W., & Li, J. (Dec. 16, 2022). Aging and aging-related diseases: from molecular mechanisms to interventions and treatments. *Nature: Signal transduction and targeted therapy*.
- Gustafsson, T., & Ulfhake, B. (2024). Aging skeletal muscles: What are the mechanisms of age-related loss of strength and muscle mass, and can we impede its development and progression? *International Journal of Molecular Science*, 25(20), 10932.
- Guttman, D.L. (1974). Alternatives to disengagement: Aging among the Highland Druze. In R. LaVine (Ed.), *Culture and personality: Contemporary readings*. Chicago: Aldine.
- Haapanen, N., Miilunpalo, S., Vuori, I., Oja, P., & Pasnen, M. (1996). Characteristics of leisure time physical activity associated with decreased risk of premature all-cause and cardiovascular disease mortality in middle-aged men. *American Journal of Epidemiology*, 143, 870-880.
- Haines, C., Chung, T., Chang, A., Masarei, J., & Tomlinson, B. (1996). Effect of oral estradiol on Lp(a) and other lipoproteins in postmenopausal women. *Archives of Internal Medicine*, 156, 866-872.
- Hagestad, G., & Neugarten, B. (1985). Age and the life course. In R.H. Binstock and E.Shanas (Eds.), *Handbook of aging and the social sciences* (2nd ed.). New York: Van Nostrand.
- Halliwell, B., & Gutteridge, J.M.C. (2015). Free radicals in biology and medicine. *Oxford University Press*, p. 896.
- Hansen, T. & Blekesaune, M. (2022). The age and well-being "paradox:" A longitudinal and multidimensional reconsideration. *European Journal of Ageing*, Vol. 19, pp. 1277-1286.
- Hansen, T. & Slagsvold, B. (2012). The age and subjective well-being paradox revisited: A multidimensional perspective. *Norsk Epidemiological*, 2, 187-195. Retrieved from <http://www.ntnu.no/ojs/index.php/norepid/article/view/1565/1459>
- Harlow, R.E., & Cantor, N. (1996). Still participating after all these years: A study of life task participation in later life. *Journal of Personality and Social Psychology*, 71, 1235-1249.
- Hayflick, L. (1994). *How and why we age*. New York: Ballantine Books.
- Hayflick, L., & Moorhead, P.S. (1961). The serial cultivation of human diploid cell strains. *Experimental Cell Research*, 25, 585-621.
- Heckhausen, J. (1997). Developmental regulation across adulthood: Primary and secondary control of age-related challenges. *Developmental Psychology*, 33, 176-187.
- Heidenreich, P.A. et al (2011) Forecasting the future of cardiovascular disease in the United States: A policy statement from the American. *Heart Association Circulation*, 123: 933–944.
- Henkel, L.A., Johnson, M.K. & De Leonardis, D.M. (1998). Aging and source monitoring: Cognitive processes and neuropsychological correlates. *Journal of Experimental Psychology: General*, 127, 251-268.
- Herrmann M., Pusceddu I., Marz W., & Herrmann W. (2018). Telomere biology and age-related diseases. *Clinical Chemistry and Laboratory Medicine*. 56:1210–1222.
- Herskind, A.M., McGue, M., Holm, N.V., Sorensen, T.L., Harvald, B., & Vaupel, J.W. (1996a). The heritability of human longevity: a population-based study of 2872 Danish twin pairs born 1870-1900. *Human Genetics*, 97, 319-323.
- Herskind, A.M., McGue, M., Iachine, I.A., Holm, N., Sorensen, T.I., Harvald, B., & Vaupel, J.W. (1996b). Untangling genetic influences on smoking, body mass index and longevity: A multivariate study of 2464 Danish twins followed for 28 years. *Human Genetics*, 98, 467-475.
- Hess, T. M., & Queen, T. L. (2014). Aging influences on judgment and decision processes: Interactions between ability and experience. In Verhaeghen, P. & Hertzog, C. (Eds.), *Emotion, social cognition, and everyday problem solving during adulthood* (pp. 238–255). New York: Oxford University Press.
- Hochschild, A. (1975). Disengagement theory: a critique and proposal. *American Sociological Review*, 40, 553-569.
- Holahan, C. (1984). Marital attitudes over 40 years: A longitudinal and cohort analysis. *Journal of Gerontology*, 39, 49-57.
- Hong, H., Lachman, M.E., Charles, S.T. et al. (2021). The positive influence of sense of control on physical, behavioral, and psychosocial health in older adults: An outcome-wide approach. *Preventive Medicine*. Vol. 149.
- Hooymann, N., & Kiyak, H.A. (1993). *Social gerontology: A multidisciplinary perspective* (3rd ed). Needham Heights, MA: Allyn and Bacon.
- Hoyert, D.L., Kochanek, K.D., & Murphy, S.L. (1999). Deaths: Final data for 1997. *National Vital Statistics Reports*, 47, No. 19.
- Hughes, M.L. & Lachman, M.E. (2018). Social comparisons of health and cognitive functioning contribute to changes in subjective age. *Journal of Gerontology B Psychological Science Social Science*. 14:73(5): pp. 816-824.
- Hui, J., He, S., Liu, R. et al. (2020). Trends in erectile dysfunction research from 2008 to 2018: a bibliometric analysis. *International Journal of Impotence Research*, Vol. 32, pp. 409–419.
- Iwasaki, T., Kimura, H., Tanaka, K. et al. (2023). Association between height loss and mortality in the general population. *Scientific Reports*, Vol. 13, p. 3593.
- Izquierdo, M., Merchant, R.A., Morley, J.E. et al. (2021). International exercise recommendations in older adults: Expert consensus guidelines. *The Journal of Nutrition, Health, & Aging*. Vol. 25, pp. 824-853.
- Jackson, J. B., Miller, R. B., Oka, M., & Henry, R. G. (2014). Gender differences in marital satisfaction: A meta-analysis. *Journal of Marriage and Family*, 76(1), 105–129.
- Jett, K. (2016). Theories of aging. In T.A. Touhy & Jett, K. (Eds.), *Ebesele & Hess Toward healthy aging* (9th ed), pp. 31-39).
- Jeżewska-Zychowicz, M. & R. Gajda, R. (2023). Relationship between the Prevalence of Metabolic Disease and Impaired Mobility, Diet, Physical Activity, and Socio-Demographic Characteristics in the Polish Elderly—A Cross-Sectional Study. *Life*, 13(4), p. 864.
- Jiang, D., Fung, H. H., Sims, T., Tsai, J. L., & Zhang, F. (2016). Limited time perspective increases the value of calm. *Emotion*, 16(1), 52–62.
- Jianping, J., Zhao, J., Liu, Z., et al. (2023). Association between healthy lifestyle and memory decline in older adults: 10 year,

- population based, prospective cohort study. *British Medical Journal*, 380.
- Jomova, K., Raptova, R., & Alomar, S.Y. et al. (2023). Reactive oxygen species, toxicity, oxidative stress, and antioxidants: chronic diseases and aging. *Archives of Toxicology*, 97, 2499-2574.
- Jones, S.A., & Noppeney. (2024). older adults preserve audiovisual integration through enhanced cortical activations, not by recruiting new regions. *PLOS Biology*, 22(2).
- Jurivich, D.A., Qui, L., & Welk, J.F. (1997). Attenuated stress responses in young and old human lymphocytes. *Mechanisms of Ageing and Development*, 94, 233-249.
- Kahn, R.L., & Antonucci, T.C. (1980). Convoys over the life course: Attachment, roles, and social support. In P.B. Baltes & O.G. Brim (Eds.), *Life span development and behavior* (Vol. 3, pp. 253-286). San Diego, CA: Academic Press.
- Kanabar, R., Mazur, A., Plum, A., & Schmied, J. (2022). Correlates of testosterone change as men age. *The Aging Male*, 25(1), pp. 29-40.
- Kane, R.C., & Kane, R.A. (1990). Health care for older people: Organizational and policy issues. In R. Binstock and L. George (Eds.), *Aging and the social sciences* (3d ed.). New York: Academic Press.
- Kannel, W., D'Agostino, R., & Cobb, J. (1996). Effect of weight on cardiovascular disease. *American Journal of Clinical Nutrition*, 63(suppl.), 419S-422S.
- Kasper, J. (1988). *Aging alone: Profile and projections*. Baltimore, MD: Commonwealth Fund Commission.
- Kaufman, A.S., Kaufman, J.L., McLean, J.E., & Reynolds, C.R. (1991). Is the pattern of intellectual growth and decline across the adult life span different for men and women? *Journal of Clinical Psychology*, 47, 801-812.
- Kay, A.C., Whitson, J.A., Galinsky, A.D. (2009). Compensatory control: achieving order through the mind, our institutions, and the heavens. *Current Dir Psychology Sci*, 18(5), pp. 264-268.
- Kelly, J. (2014). *Biology of Aging*. Lumen Learning.
- Kemper, S. (1992). Language and aging. In F.I.M. Craik & T.A. Salthouse (Eds.), *The handbook of aging and cognition* (pp. 213-270). Hillsdale, NJ: Lawrence Erlbaum.
- Kim, Y.J., & Lee, C.S. (2015). Effects of grit on the successful aging of the elderly in Korea. *Indian Journal of Science and Technology*, 8(57), 373-378.
- Kim, J.C., Park, T.J., & Kang, H.Y. (2022). Skin-aging pigmentation: Who is the real enemy? *Cells*, 11(16), 2541.
- Klein, S., Gastadelli, A., Yki-Jarvinen, H., & Scherer, P.E. (January 4, 2022). *Cell Metabolism*, Vol. 34, Issue 1, pp. 11-20.
- Knopman, D. S. et al. (2012). Alzheimer disease. *Nat. Rev. Dis. Prim.* 7, 33.
- Koehly, L.M., & Manalel, J.A. (June, 2023). Interconnected social convoys: Understanding health and well-being through linked personal networks. *Advances in Life Course Research*, Vol. 56. Elsevier.
- Korczyn, A.D., & Grinberg, L.T. (2024). Is Alzheimer disease a disease? *Nature Reviews Neurology*, Vol. 20, 245-251.
- Kotler-Cope, S., & Camp, C. (1990). Memory interventions in aging populations. In E.A. Lovelace (Ed.), *Aging and cognition: Mental processes, self-awareness and interventions* (pp. 231-261). Amsterdam: North Holland.
- Kozbelt, A. (2008). Longitudinal hit ratios of classical composers: Reconciling "Darwinian" and expertise acquisition perspectives on lifespan creativity. *Psychology of Aesthetics, Creativity and the Arts*, 2(4), 221-235.
- Kramarow, E., Lentzner, H., Rooks, R., Weeks, J., & Saydah, S. (1999). Health and Aging Chartbook. *Health, United States, 1999*. Hyattsville, MD: National Center for Health Statistics.
- Kwiek, M., & Szymula, L. (2025). Quantify lifetime productivity changes: A longitudinal study of 320,000 late-career scientists. *Quantitative Science Studies, Advance Publication*. <https://doi.org/10.1162/qss.a.16>
- Labouvie-Vief, G., Hakim-Larson, J., & Hobart, C.J. (1987). Age, ego level, and the life-span development of coping and defense processes. *Psychology and Aging*, 2, 286-293.
- Lachman, M.E., & Weaver, S.L. (1998). Sociodemographic variations in the sense of control by domain: Findings from the McArthur studies of midlife. *Psychology and Aging*, 13, 553-562.
- Lachman, M.E., & Weaver, S.L., Bandura, M., Elliott, E., & Lewkowicz, C.J. (1992). Improving memory and control beliefs through cognitive restructuring and self-generated strategies. *Journal of Gerontology: Psychological Sciences*, 47, P293-299.
- Larrabee, G.J. (November 30, 2019). *Memory changes in the aging brain*. Cambridge University Press.
- Launer, L.J., Masaki, K., Petrovitch, H., Foley, D., & Havlik, R.J. (1995). The association between midlife blood pressure levels and late-life cognitive function. The Honolulu-Asia Aging Study. *Journal of the American Medical Association*, 274, 1846-1851.
- Lawton, M.P., Kleban, M.H., Rajagopal, D., & Dean, J. (1992). Dimensions of affective experience in three age groups. *Psychology and Aging*, 7, 171-184.
- Lawton, M.P., & Nahemow, L. (1973). Ecology and the aging process. In C. Eisdorfer & M.P. Lawton (Eds.), *The psychology of adult development and aging*. Washington, DC: American Psychological Association.
- Lee, P.R. & Benjamin, A.E. (1989). Health policy trends: Impact on the academic health center. In S. Andreopoulos and J. Hogness (Eds.), *Health care for an aging society*. New York: Churchill Livingstone.
- Lehman, H.C. (1953). *Age and achievement*. Princeton, NJ: Princeton University Press.
- Lemon, B., Bengston, V., & Peterson, J. (1972). Activity types and life satisfaction in a retirement community. *Journal of Gerontology*, 27, 511-523.
- Li, Y., Baldassi, M., Johnson, E. J., & Weber, E. U. (2013). Complementary cognitive capabilities, economic decision making, and aging. *Psychology and Aging*, 28, 595-613.
- Li, Y., Gao, J., Enkavi, A. Z., et al. (2015). Sound credit scores and financial decisions despite cognitive aging. *Proceedings of the National Academy of Sciences USA*, 112, 65-69
- Li, A., Zhang, Z, Ren, Y., Wang, Y., Fang, J., Yue, H., Ma, S., & Guan, F. (2021). Aging and age-related diseases: from mechanisms to therapeutic strategies. *BioGerontology*, 22: 165-187.
- Löckenhoff, C. E., & Carstensen, L. L. (2007). Aging, emotion, and health-related decision strategies: Motivational manipulations can reduce age differences. *Psychology and Aging*, 22, 134-146
- Longino, C.F., & Kart, C.S. (1982). Explicating activity theory: A formal replication. *Journal of Gerontology*, 35, 713-722.
- Lopez-Lluch, G., Navas, P. (2016). Calorie restriction as an intervention in ageing. *The Journal of Physiology*. Wiley Online Library.
- Lopata, H. (1979). *Women and widows*. New York: Elsevier.
- Longino, C.F., & Kart, C.S. (1982). Explicating activity theory: A formal replication. *Journal of Gerontology*, 35, 713-722.
- Lopez-Lluch, G., Navas, P. (2016). Calorie restriction as an intervention in ageing. *The Journal of Physiology*. Wiley Online Library.
- Lueckenotte, A.G. (2000). *Theories of Aging. Gerontologic Nursing*. St. Louis: Mosby.
- MacKay, D.G., & Abrams, L. (1996). Language, memory and aging: Distributed deficits and the structure of new-old connections. In J.E. Birren, K.W. Schaie, R.P. Abeles, M. Gatz,

- & T.A. Salthouse (Eds.), *Handbook of the psychology of aging* (4th ed.) (pp. 251-265). San Diego, CA: Academic Press.
- Manalel, J.A. & Antonucci, T.C. (Sep. 2022). Development of social convoys: Trajectories of convoy structure and composition from childhood through adulthood. *Developmental Psychology*, 58(9):1806-1815.
- Marieb, E.N., & Hoehn, K. (2019). *Human anatomy and physiology*, 9th edition.
- Marshall, V.W. (1996). The stage of theory in aging and the social sciences. In R.H. Binstock & L.K. George (eds.), *Handbook of aging and the social sciences* (4th ed. pp. 12-26). San Diego, CA: Academic Press.
- Marshall, V.W., & Clarke, P.J. (2007). *Theories of Aging: Social*. Elsevier Inc.
- Marsiglio, W., & Donnelly, D. (1991). Sexual relations in later life: A national study of married persons. *Journal of Gerontology: Social Sciences*, 46, 338-344.
- Masters, W.H., & Hognson, V.E. (1970). *Human sexual inadequacy*. Boston: Little, Brown.
- Masters, W.H., & Johnson, V.E. (1966). *Human sexual response*. Boston: Little, Brown.
- Mata, R., Pachur, T., von Helversen, B., et al. (2012). Ecological rationality: Framework for understanding and aiding the aging decision maker. *Frontiers in Neuroscience*, 6, 19.
- Mazzilli, F. (2022). Erectile Dysfunction: Causes, diagnosis and treatment: An update. *Journal of Clinical Medicine*, 11(21), 6429.
- McCrae, R.R., & Costa, P.T., Jr. (1990). *Personality in adulthood*. New York: Guilford.
- McDarby, M., Kozlov, E.K., & Carpenter, B. (Nov. 8, 2019). Frequency of contact among family members and end-of-life care planning. *Innovation in Aging*, Vol. 3 (Supplement 1): S669-670.
- Mensa, E., Latini, S., Ramini, D., Storci, G., Bonafe, M., & Olivieri, F. (2019). The telomere world and aging: analytical challenges and future perspectives. *Ageing Research Reviews*, 50:27-42
- Meyer, B.J.F., Russo, C., & Talbot, A. (1995). Discourse comprehension and problem solving: Decisions about the treatment of breast cancer by women across the life span. *Psychology and Aging*, 10, 84-103.
- Michalos, A.C. (1985). Multiple discrepancies theory (MDT). *Social Indicators Research*, 16, 347-413.
- Mikels, J. A., Löckenhoff, C. E., Maglio, S. J., et al. (2010). Following your heart or your head: Focusing on emotions versus information differentially influences the decisions of younger and older adults. *Journal of Experimental Psychology: Applied*, 16, 87-95.
- Minkler, M., & Estes, C. (1984). *Readings in the political economy of aging*. Farmingdale, N.Y.: Baywood.
- Monfared, A.A.T., Hummel, N., Chandak, A., Khachatryan, A., Zhang, R., & Zhang, Q. (2024). Prevalence estimation of Dementia/Alzheimer's Disease using health and retirement study database in the United States. *The Journal of Prevention of Alzheimer's Disease*, Vol. 11, Issue 5, pp. 1183-1188.
- Moqri, M., Herzog, C., Poganik, J., et al. (2023). Biomarkers of aging for the identification of longevity interventions. *Cell*. Vol. 186, Issue 18, pp. 3758-3775.
- Mroczek, D.K., & Kolarz, C.M. (1998). The effect of age on positive and negative affect: A developmental perspective on happiness. *Journal of Personality and Social Psychology*, 75, 1333-1349.
- Multhaup, K.S., Balota, D.A., & Cowan, N. (1996). Implications of aging, lexicality, and item length for the mechanisms underlying memory span. *Psychonomic Bulletin and Review*, 3, 112-120.
- Muraki, K., Nyhan, K., Han, L., Murnane, J.P. (2012). Mechanisms of telomere loss and their consequences for chromosome instability. *Frontiers in Oncology*, 2:135.
- Naso, C.M., Sy, L., Song, G., et al. (2025). Time trend analysis of osteoporosis prevalence among adults 50 years of age and older in the USA, 2005-2018. *Osteoporosis International*, 36, 547-554.
- National Institute of Diabetes and Kidney Disease (1999). Impotence. [www.niddk.nih.gov/health/urolog/pubs/impotence/impotence.htm](http://www.niddk.nih.gov/health/urolog/pubs/impotence/impotence.htm).
- National Institutes of Health (1992). *Impotence*. National Institutes of Health consensus development conference statement. Bethesda, MD: National Institutes of Health.
- Nelson, E.A., & Dannefer, D. (1992). Aged heterogeneity: Fact or fiction? The fate of diversity in gerontological research. *Gerontologist*, 32, 17-23.
- Neugarten, B., Havighurst, R.J., & Tobin, S.S. (1968). In B.L. Neugarten (Ed.), *Personality and pattern of aging in middle age and aging*. Chicago: University of Chicago Press, 173-177.
- Neumann, C.V. (2000). *Sources of meaning and energy in the chronically ill frail elder*. Unpublished paper prepared for the Ronald E. McNair Research Program, University of Wisconsin-Milwaukee.
- NIH (National Institute of Health). (2020). *Is longevity determined by genetics?* <https://ghr.nlm.nih.gov/primer/traits/longevity>
- Orgel, L.E. (1963). The maintenance of the accuracy of protein synthesis and its relevance to aging. *Proceedings of the National Academy of Science of the USA*, 49, 512-517.
- Overstreet, D.S., Strath, L.J., Jordan, M.K., Jordan, I., Hobson, J.M., Owens, M.A., Williams, A.C., Edwards, R.R., & Meints, S.M. (2023). A brief overview: Sex differences in prevalent chronic musculoskeletal conditions. *International Journal of Environmental Research and Public Health*, 20(5), 4521.
- Paffenbarger, R.S., Hyde, R.T., Wing, A.L., Lee, I.M., Jung, D.L., & Kampert, J.B. (1993). The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *New England Journal of Medicine*, 328, 538-545.
- Park, D.C., & Reuter-Lorenz, P. (2009). The adaptive brain: aging and neurocognitive scaffolding. *Annual Review of Psychology*, 60, 173-196.
- Park, D.C., Smith, A.D., & Cavanaugh, J.C. (1990). Metamemories of memory researchers. *Memory and Cognition*, 18, 321-327.
- Peelle, J.E. (2019). Language and aging In Greig I. de Zubizaray & Niels O. Schiller (eds.), *The Oxford Handbook of Neurolinguistics*, 295-316.. Oxford: Oxford University Press.
- Perkins M.M., Ball M.M., Whittington F.J., & Hollingsworth C. (April 1, 2012). Relational Autonomy in Assisted Living: A Focus on Diverse Care Settings for Older Adults. *Journal of Aging Studies*. 26(2):214-225.
- Perrig, W.J., Perrig, P., & Stahelin, H.B. (1997). The relation between antioxidants and memory performance in the old and very old. *Journal of the American Geriatrics Society*, 45, 718-724.
- Pfeiffer, E., Verwoerd, A., & Davis, G.C. (1974). Sexual behavior in aged men and women. In E. Palmore (Ed.), *Normal Aging II* (pp. 243-251). :Durham, NC: Duke University Press.
- Portin, R., Saarijaervi, S., Joukamaa, M., & Salokangas, R.K.R. (1995). Education, gender and cognitive performance in a 62 year-old normal population: Results from the Turva Project. *Psychological Medicine*, 25, 1295-1298.
- Pradipta, A.R., Mariana, M., & Idris, F. (2025). A cortisol dynamics in the elderly: Systematic Review. *Health System and Medical Journal*, Vol. 1, No. 1.
- Punyiczki, M., & Fesus, L. (1998). Heat shock and apoptosis. In P. Csermeley (Ed.), *Stress of life: From molecules to man*. *Annals of the New York Academy of Sciences*, 851, 67-74.
- Reuter-Lorenz, P.A., & Cappell, K.A. (2008). Neurocognitive aging and the compensation hypothesis. *Current Directions in Psychological Science*, 17(3), 177-182.
- Reuter-Lorenz, P.A., & Lustig, C. (2005). Brain aging: Reorganizing discoveries about the aging mind. *Current Opinion in Neurobiology* 15(2). 245-251.

## HEALTHY AGING

- Reuter-Lorenz, P.A., & Park, D.C. (2014). How does it STAC Up? Revisiting the Scaffolding Theory of Aging and Cognition. *Neuropsychology Review* 24, 355-0370.
- Rhodes, E., & Giovannetti, T. (2017). Grit and successful aging in older adults. Presented at the 45th Annual Meeting of the International Neuropsychological Society, New Orleans, LA, February.
- Rhodes, E., & Giovannetti, T. (May 28, 2020). Grit and successful aging. In part IV - Cognitive, Social, and Biological Factors Across the Lifespan. (Eds.), AyannaThomas & Angela Gutches. *Cambridge University Press*.
- Ricklefs, R.E., & Finch, C.E. (1995). *Aging: A natural history*. New York: Scientific American Library.
- Riley, M.W. (1994). Age integration and the lives of older people. *Gerontologist*, 34, 110-115.
- Riley, M.W., Johnson, J., & Forner, A. *Aging and society* (Vol.3): *A sociology of age stratification*. New York: Russell Sage Foundation.
- Rimm, E., Ascherio, A., Giovannucci, E., Spiegelman, D., & Stampfer, M. (1996). High-fiber diet can reduce heart attack risk. *Journal of the American Medical Association*, 275, 447-451.
- Robine, J., & Vaupel, J.W. (2001). Supercentenarians: slower aging individuals or senile elderly? *Experimental Gerontology*, 36, (4-6), 915-930.
- Rosenberg, E. (2022). Social theories of aging. *Europasian Journal of Medical Sciences*, 4:20-30.
- Rosow, I. (1985). Status and role change through the life cycle. In R.H. Binstock and E. Shanas (Eds.), *Handbook of aging and the social sciences* (2nd ed.). New York: Van Nostrand, 62-93.
- Rothbaum, F., Weisz, J.R., & Snyder, S.S. (1982). Changing the world and changing the self: A two process model of perceived control. *Journal of Personality and Social Psychology*, 42, 5-37.
- Rowe, J.W. (2018). A research network to support a productive, equitable, aging society. McArthur Foundation.
- Rowe, J.W., & Kahn, R.L. (1998). *Successful aging*. New York: Pantheon Books.
- Sadaqa, M., Nemeth, Z., Makai, A. et al. (August 2, 2023). Effectiveness of exercise interventions on fall prevention in ambulatory community-dwelling older adults: A systematic review with narrative synthesis. *Frontiers in Public Health*, Vol. 11.
- Salthouse, T.A. (1985). Speed of behavior and its implications for cognition. In J.E. Birren & K.E. Schaie (Eds.), *Handbook of the psychology of aging* (2nd ed.) (pp. 400-426). New York: Van Nostrand Reinhold.
- Salthouse, T.A. (2010). Major issues in cognitive aging. *Oxford University Press*. New York.
- Salthouse, T.A., & Babcock, R.L. (1991). Decomposing adult age differences in working memory. *Developmental Psychology*, 27, 763-776.
- Sattaur, Z., Lashley, L., Golden, C. (April 25, 2025). Wear and tear theory of aging. *Essays in Developmental Psychology*. Nova Southeastern University.
- Sattaur, Z., Lashley, L., Golden, C. (April 25, 2025). Wear and tear theory of aging. *Essays in Developmental Psychology*. Nova Southeastern University.
- Saxon, S., Eiten, M.J., Perkins, E.A. (2022). *Physical change and aging: A guide for the helping professions*. (7th edition). Springer Publishing Company.
- Schaie, K.W. (1983). The Seattle Longitudinal Study: A 21-year exploration of psychometric intelligence in adulthood. In K.W. Schaie (Ed.), *Longitudinal studies of adult psychological development* (pp. 64-135). New York: Guilford.
- Schaie, K.W. (1996). Intellectual development in adulthood. In J.R. Birren, K.W. Schaie, R.P. Abeles, M. Gatz, & T.A. Salthouse (Eds.), *Handbook of the psychology of aging* (4th ed.) (pp. 266-286). San Diego CA: Academic Press.
- Schaie, K. Warner. (2005). Developmental Influences on Adult Intelligence: *The Seattle Longitudinal Study*. Oxford University Press.
- Schaie, K.W., Dutta, R., & Willis, S.L. (1991). Relationship between rigidity-flexibility and cognitive abilities in adulthood. *Psychology and Aging*, 6, 371-383.
- Schneider, J.L., Rowe, J.H., Garcia-de-Alba, C. (2021). The aging lung: Physiology, disease, and immunity. *Cell*, Vol. 184, Issue 8, pp. 1990-2019.
- Schoenhofen, F.A., Wyszynski, D.F., Andersen, S., et al. (2006). Characteristics of 32 supercentenarians. *JAGS*, 54, 1237-1240.
- Schroots, J.J.F. (1996). Theoretical developments in the psychology of aging. *Gerontologist*, 36, 742-748.
- Schumacher, B., Pothof, J., Vijg, J., & Hoeijmakers, H.J. (2021). The central role of DNA damage in the ageing process. *Nature*, 592, pp. 695-703.
- Sebastiani, P., Fangui, X.S., Andersen, S., et al. (2013). Families enriched for exceptional longevity also have increased health-span: Findings from the Long Life Family Study. *Front Public Health*, 1, 38.
- Sellami, M., Bragazzi, N.L., Aborghaba, B., & Elrayess, M.A. (April 13, 2021). The impact of acute and chronic exercise on immunoglobulins and cytokines in elderly: Insights from a critical review of the literature. *Frontiers in Immunology*, Vol. 12.
- Sen, M. (2025). Treating ageing: A review. *Research Journal of Life Sciences, Bioinformatics Pharmaceutical and Chemical Sciences*. <http://www.rjlbpcs.com>
- Shadden, B.B. (1997). Discourse behaviors in older adults. *Seminars in Speech and Language*, 18, 143-156.
- Shafto, M.A., & Tyler, L.K. (2014). Language in the aging brain: The network dynamics of cognitive decline and preservation. *Science* 346 (6209). 583-587.
- Shah, S.J., Fang, M.C., Wannier, S.R. et al. (2022). Association of social support with functional outcomes in older adults who live alone. *JAMA Internal Medicine*, Vol.182(1): pp. 26-32.
- Shanas, E. (1979). The family as social support in old age. *The Gerontologist*, 19, 169-174.
- Shanas, E. (1980). Older people and their families: The new pioneers. *Journal of Marriage and the Family*, 42, 9-14.
- Shock, N.W. (1985). Longitudinal studies of aging in humans. In C.E. Finch & E.L. Schneider (Eds.), *Handbook of the biology of aging* (2nd ed.) (pp. 721-743). New York: Van Nostrand.
- Simonton, D.K. (1998). Career paths and creative lives: A theoretical perspective on late-life potential. In C. Adams-Price (Ed.), *Creativity and successful aging: Theoretical and empirical approaches* (pp. 3-18). New York: Springer.
- Singh, M.K., Shin, Y., Ju, S., et al. (2024). Heat shock response and heat shock proteins: Current understanding and future opportunities in human diseases. *International Journal of Molecular Science*, 25(8), p. 4200.
- Sinnott, J.D. (1989). A model for solution of ill-structured problems: Implications for everyday and abstract problem-solving. In J.D. Sinnott (Ed.), *Everyday problem solving: Theory and applications* (pp. 72-99). New York: Praeger.
- Smith, A.D. (1996). Memory. In J.E. Birren, K.W. Schaie, R.P. Abeles, M. Gatz, & T.A. Salthouse (Eds.), *Handbook of the psychology of aging* (4th ed.) (pp. 236-250). San Diego, CA: Academic Press.
- Smith, J.R., & Pereira-Smith, O.M. (1996). Replicative senescence: Implications for in vivo aging and tumor suppression. *Science*, 273, 63-67.
- Smith, B.D., Smith, G.L., Hurria, A., Hortobagyi,

- G.N., Buchholz, T.A. (2009). Future of cancer incidence in the United States: burdens upon an aging changing nation. *Journal of Clinical Oncology*, 27:2758–2765
- Snowden, D.A. (1997). Aging and Alzheimer's disease: Lessons from the Nun Study. *Gerontologist*, 37, 150-156.
- Sohal, R.S., & Weindruch, R. (1996). Oxidative stress, caloric restriction, and aging. *Science*, 273, 59-63.
- Sonkin, D., Thomas, A., & Teicher, B.A. (2024). Cancer treatments: past, present, and future. *Cancer Genetics*, Vol. 286-287, pp. 18-24.
- Starr, B.D. (1985). Sexuality and aging. In M.P. Lawton & G.L. Maddox (Eds.), *Annual Review of Gerontology and Geriatrics* (Vol 5, pp. 97-126). New York: Springer.
- Starr, B.D., & Weiner, M.B. (1981). *The Starr-Weiner report on sex and sexuality in the mature years*. New York: McGraw-Hill.
- Stephens, C., Breheny, M. (2018). Healthy Aging: A capability approach to inclusive policy and practice. Chapter 6, *Social Connections*. Taylor and Francis.
- Stevens, J., Cai, J., Pamuk, E.R., Williamson, D.F., Thun, M.J., & Wood, J.L. (1998). The effect of age on the association between body-mass index and mortality. *New England Journal of Medicine*, 338, 1-7.
- Stewart, C. C., Yu, L., Wilson, R. S., Bennett, D. A., & Boyle, P. A. (2018). Correlates of healthcare and financial decision making among older adults without dementia. *Health Psychology*, 37, 618–626.
- Stoller, E.P. (1990). Males as helpers: The roles of sons, relatives and friends. *The Gerontologist*, 30, 228-235.
- Strough, J., Bruine de Bruin, W., & Parker, A. M. (2018). Taking the biggest first: Age differences in preferences for monetary and hedonic sequences. *Journals of Gerontology, Series B: Psychological Sciences and Social Sciences*, 74(6), 964–974
- Strough, J., Bruine,de Bruin, W.B., & Wilson, J.M. (2020). Age differences in decision making. In the *Cambridge Handbook of Cognitive Aging: A life course perspective* (Eds.), Thomas, A.K., & Gutchess, A. Cambridge University Press. Chapter 12, Part II.
- Strough, J., Mehta, C. M., McFall, J. P., & Schuller, K. L. (2008). Are older adults less subject to the sunk-cost fallacy than younger adults? *Psychological Science*, 19, 650–652.
- Strough, J., Parker, A., & Bruine de Bruin, W. (2019). Thinking about the future reduces failure to act after a missed opportunity. *Psychology and Aging*, 34(2), 311–316.
- Suominen, H. (1997). Changes in physical characteristics and body composition during 5-year follow-up in 75- and 80-year-old men and women. *Scandinavian Journal of Social Medicine. Supplementum*, 53, 19-24.
- Swift, H.J., Vauclair, M.C., Abrams, D., et al. (2014). Revisiting the paradox of well-being: The importance of national context. *The Journal of Gerontology: Series B*, Vol. 69, Issue 6, pp. 920-929.
- Taffert, G.E. (1996). Age-related physiologic changes. In D.B. Reuben, T.T. Yoshikawa, & R.W. Besdine (Eds.), *Geriatrics review syllabus* (3rd ed., pp. 11-15). Dubuque, IA: Kendall/Hunt.
- Taffert, G.E. (2023). Physiology of Aging. In: Wasserman, M.R., Bakerjian, D., Linnebur, S. et al. (eds.). (2023). *Geriatric Medicine*. Springer, Cham.
- Tenchov R., Sasso, J.M., Wang, X., & Qionggiong, A.Z., (2023). Aging hallmarks and progression and age-related diseases: A landscape view of research advancement. *ACS Chemical Neuroscience*, Vol. 1, Issue 15.
- Teodoro, T.H., Costa, K.P.M., Prestes, J., et al. (February, 2025) The effects of acute and chronic exercises on immune markers of TH1/TH2 cells in older adults: A systematic review. *Frontiers in Physiology*, Vol. 16.
- Thomas, C., Rukhsana, G., Mirza, G., et al. (2021). Age-related macular degeneration. *Medical Clinics of North America*. Vol. 105, Issue 3, pp. 473-491.
- Thornton, R., & Light, L.L. (2006). Language comprehension and production in normal aging. In K. Warner Schaie & Sherry L Willis (eds.), *Handbook of the psychology of aging*, 261-287. Amsterdam: Academic Press.
- Thurstone, L.L. (1938). *Primary mental abilities*. Chicago: University of Chicago Press.
- Tinsdale, J.A., & Marshall, V.W. (1980). A generational conflict perspective for gerontology. In V.W. Marshall (Ed.), *Aging in Canada: Social perspectives*. Don Mills, Ontario: Fitzhenry and Whiteside, 43-50.
- Touhy, T.A., & Jett, K. (2022). *Ebersole and Hess' Gerontological Nursing & Healthy Aging* (6th ed.). Elsevier.
- Troll, L.E., & Skaff, M.M. (1997). Perceived continuity of self in very old age. *Psychology and Aging*, 12(1), 162-169.
- Trott, C.T., Friedman, D., Ritter, W., & Fabiani, M. (1997). Item and source memory: Differential age effects revealed by event-related potentials. *Neuroreport*, 8, 3373-3378.
- Tsvetanov, K.A, Mevorach, C., & Humphreys, G.W. (2013). Age-related differences in selection by visual saliency. *Attention Perception Psychophysiology*, 75, 1382-1394.
- Uchino, B.N., Cacioppo, J.T., & Kiecolt-Glaser, J.K. (1996). The relationship between social support and physiological processes. A review with emphasis on underlying mechanisms and implications for health. *Psychological Bulletin*, 119, 488-531.
- Uchino, B.N., Trettevik, R., Kent, D., Grey, R.B., et al. (2018). Social support, social integration, and inflammatory cytokines: A meta-analysis. *Health Psychology*, 37(5), 462-471.
- Uhlenberg, P. (1996). The burden of aging: A theoretical framework for understanding the shifting balance of care giving and care receiving as cohorts age. *Gerontologist*, 36, 761-767.
- Uhlenberg, P. (2000). Why study age integration? *Gerontologist*, 40, 261-266.
- U.S. Senate Special Committee on Aging. (1986). *Developments in aging: 1985*. Washington, D.C.: U.S. Government Printing Office.
- U.S. Senate Special Committee on Aging. (1992). *Aging America: Trends and projections, 1991 edition*. Washington, D.C.: U.S. Department of Health and Human Services.
- U.S. Senate Special Committee on aging. (1992). *Aging America: Trends and projections*. Washington, D.C.: U.S. Government Printing Office.
- Vaiserman, A., Krasnienkov, D. (2021). Telomere length as a marker of biological age: State-of-the-art, open issues, and future perspectives. *Frontiers in Genetics*, January 21.
- van Boxtel, M.P., Paas, F.G., Houx, P.J., Adam, J.J., Teeken, J.C., & Jolles, J. (1997). Aerobic capacity and cognitive performance in a cross-sectional aging study. *Medicine and Science in Sports and Exercise*, 29, 1357-1365.
- Verhaeghen, P., Marcoen, A., & Goossens, L. (1993). Facts and fiction about memory aging: A quantitative integration of research findings. *Journal of Gerontology: Psychological Sciences*, 48, P157-171.
- Verhoef, P.K, Stampfer, M., Buring, J., Gaziano, J.M., & Allen, R. (1996). Homocysteine metabolism and risk of myocardial infarction: Relation with Vitamins B6, B12, and folate. *American Journal of Epidemiology*, 143, 845-859.
- Vignesawarn, K., & Hamoda, H. (2022). Hormone replacement therapy - Current recommendations. *Best Practice & Research Clinical Obstetrics & Gynaecology*, Vol. 81, pp. 8-21.
- von Zglincki, T., Nilsson, E., Docke, W.D., & Brunk, U.T. (1995). Lipofuscin accumulation and aging of fibroblasts. *Gerontology*, 41, (Suppl. 2), 95-108.

## HEALTHY AGING

- Waldo, D., Sonnefeld, S., McKusick, D., & Arnett, R. (1989). Health expenditures by age group, 1977 and 1987. *Health Care Financing Review*, 10.
- Walker, A. (1981). Toward a political economy of old age. *Aging and Society*, 1, 73-94.
- Wallston, B., Alagna, S., DeVellis, B., & DeVellis, R. (1983). Social support and physical health. *Health Psychology*, 2, 367-391.
- Wang, Y., Li, G., Yang, L., Luo, R., Rifang, L., & Guo, G. (June 2, 2022). Development of Innovative biomaterials and devices for the treatment of cardiovascular diseases. *Advanced Materials*.
- Weindruch, R., & Walford, R.L. (1988). *The retardation of aging and disease by dietary restriction*. Springfield, IL: Charles C. Thomas.
- Weiss, D., Couto, M.C.P., & Rothermund, K. (June 23, 2025). How social and temporal comparisons shape subjective aging. *Psychology of Aging*.
- Whitbourne, S.K. (2001). *Adult development and aging: Biopsychosocial perspectives*. New York: John Wiley & Sons, Inc.
- Wickramasinghe, K., Mathers, S., Wopereis, D.S. & Griffiths, J.C. (2020). From lifespan to healthspan: The role of nutrition in healthy ageing. *Journal of Nutritional Science*, 9, pp. 1-10.
- Wilkniss, S.M., Jones, M.G., Korol, D.L., Gold, P.E., & Manning, C.A. (1997). Age-related differences in an ecologically based study of route learning. *Psychology and Aging*, 12, 372-375.
- Willcox, B.J., Willcox, D.C., & Ferrucci, L. (2008). Secrets of healthy aging and longevity from exceptional survivors around the globe: Lessons from octogenarians to supercentenarians. *Journal of Gerontology A Biological Science Medical Science*, 63(11), 1201-1208.
- Willis, S.L. (1990). Current issues in cognitive training research. In E.A. Lovelace (Ed.), *Aging and cognition: Mental processes, self-awareness and interventions* (pp. 263-280). Amsterdam: North Holland.
- Willis, S.L. (1996). Everyday problem solving. In J.E. Birren, K.W. Schaie, R.P. Abeles, M. Gatz, & T.A. Salthouse (Eds), *Handbook of the psychology of aging* (4th ed.) (pp. 287-307). San Diego, CA: Academic Press.
- Wingfield, A. & Grossman, M. (2006). Language and the aging brain: patterns of neural compensation revealed by functional brain imaging. *Journal of Neurophysiology* 96(6). 2830-2839.
- Wisniewski, H.M., Wegiel, J., & Kotula, L. (1996). Some neuropathological aspects of Alzheimer's disease and its relevance to other disciplines. *Neuropathology and Applied Neurobiology*, 22, 3-11.
- World Health Organization (1997). *Composite International Diagnostic Interview (CIDI)*. Geneva, Switzerland: World Health Organization.
- Wright, N.C., Looker, A.C., Saag, K.G., et al. (2014). The recent prevalence of osteoporosis and low bone mass in the United States based on bone mineral density at the femoral neck or lumbar spine. *Journal of Bone Mineral Research*, 29 (11), 2520-2526.
- Xing, Y., Xuan, F., Wang, K., & Zhang, H. (August 1, 2023). Aging under endocrine hormone regulation. *Frontiers in Endocrinology*. Vol. 23.
- Yaeger, J.J. (2019). Integumentary function. In S.F. Meiner & J.J. Yaeger (Eds.), *Gerontologic Nursing* (6th ed., pp. 279-310).
- Yang, J.H., Lee, H.C., & Wei, Y.H. (1995). Photoageing-associated mitochondrial DNA length mutations in human skin. *Archives of Dermatological Research*, 287, 641-648.
- Yang, W., Zhao, X., Renjie, C., & Fan, J. (August 29, 2023). Progress on mechanisms of age-related hearing loss. *Frontiers in Neuroscience*, Vol. 17.
- Yin, P., & Lai, K.H. (1983). A reconceptualization of age stratification in China. *Journal of Gerontology*, 38, 608-613.
- Zarit, S.H., & Zarit, J.M. (1998). *Mental disorders in older adults: Fundamentals of assessment and treatment*. New York: Guilford.
- Zhu Y., Liu X., Ding X., Wang F., & Geng X. (2019). Telomere and its role in the aging pathways: telomere shortening cell senescence mitochondria dysfunction. *Biogerontology* 20:1-16.

## TEST - HEALTHY AGING

### 7 Continuing Education Contact Hours

Click the link “New York LMSW/LCSW/LMHC/LMFT/ Psychologist Answer Sheet” on Home Page, then follow the prompts, which include making your payment, transferring your test answers to the online answer sheet, and printing your certificate immediately.

Passing is 70% or better.

### TRUE/FALSE

1. **Programmed aging and random error theories comprise the biological theories of aging.**  
A) True B) False
2. **The free radical theory is being explored as a possible cause of aging.**  
A) True B) False
3. **The majority of older people have positive and accepting attitudes toward sexuality.**  
A) True B) False
4. **Exercise can improve immune responsiveness in elder adults.**  
A) True B) False
5. **Life expectancy has fallen over the past 60 years.**  
A) True B) False
6. **Current research indicates that the aging brain maintains much of its function and structure.**  
A) True B) False
7. **Secondary aging involves later life changes due to disease.**  
A) True B) False
8. **Higher education and continued mental activity throughout life may protect against Alzheimer’s disease.**  
A) True B) False
9. **Most studies on working memory span indicate age-related deficits.**  
A) True B) False
10. **The Baltimore Longitudinal Study of Aging concludes that lifestyle decisions cannot affect the occurrence or progression of some age-related diseases.**  
A) True B) False
11. **Age-related changes in body composition, such as loss of bone mineral content, fat increase, and muscle mass decrease are related to the \_\_\_\_\_.**  
A) endocrine system  
B) respiratory system  
C) urinary system  
D) reproductive system
12. **Brain research indicates that \_\_\_\_\_.**  
A) mental stimulation can compensate for loss of neurons.  
B) remaining neurons do not increase their synapses.  
C) there is no neurological basis for memory changes in later adulthood.  
D) older adults cannot compensate for brain deficits.
13. **The leading cause of death in people over age 65 involves the \_\_\_\_\_.**  
A) reproductive system  
B) cardiovascular system  
C) vestibular system  
D) bones
14. **The formation of plaques and tangles is associated with \_\_\_\_\_.**  
A) diabetes  
B) skin cancer  
C) Alzheimer’s disease  
D) hearing impairment
15. **Sedentary lifestyle, smoking, body weight, and alcohol intake represent the four major risk factors for \_\_\_\_\_.**  
A) skin disorders  
B) diabetes  
C) heart disease  
D) Alzheimer’s disease

Continuing Psychology Education Inc. is recognized by the New York State Education Department’s State Board for: Social Work as an approved provider of continuing education for licensed social workers #SW-0387; Mental Health Practitioners as an approved provider of continuing education for licensed mental health counselors #MHC-0080, and licensed marriage and family therapists #MFT-0043; Psychology as an approved provider of continuing education for licensed psychologists #PSY-0006.

## HEALTHY AGING

16. **Older adults demonstrating more flexibility in attitude and personality style are less likely to experience a decline in \_\_\_\_\_.**
- A) endocrine system functioning
  - B) intellectual functioning
  - C) visual acuity
  - D) cardiovascular system functioning
17. **Decreasing the range of one's relationships to maximize social and emotional gains and minimize risks, illustrates \_\_\_\_\_.**
- A) age-related memory deficits
  - B) socioemotional selectivity theory
  - C) Alzheimer's disease
  - D) cognitive impairment
18. **Older adults are often \_\_\_\_\_.**
- A) not able to control expression of their emotions
  - B) afflicted by chronic negative emotions
  - C) better able to control expression of their emotions and they have fewer negative emotions
  - D) incapable of regulating their affective experiences
19. **Older people living alone without social support systems, compared to those with strong social supports, have greater likelihood of \_\_\_\_\_.**
- A) institutionalization
  - B) lower personally reported well-being
  - C) more difficulty adjusting to widowhood
  - D) all of the above
20. **The concepts of increased feelings of satisfaction and positive view of self, with aging, despite hardships, \_\_\_\_\_.**
- A) rarely occurs
  - B) is an unrealistic expectation
  - C) are overly dependent upon financial status
  - D) seems to be the norm

Log on to [www.texcpe.com](http://www.texcpe.com) and click the link entitled "New York LMSW/LCSW/LMHC/LMFT/Psychologist Answer Sheet" on Home Page, then follow the prompts which include making your payment, transferring your test answers to the online answer sheet, and printing your certificate immediately.